

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
12 May 2005 (12.05.2005)

PCT

(10) International Publication Number  
**WO 2005/042554 A1**

(51) International Patent Classification<sup>7</sup>: **C07H 17/00**,  
17/08, A61K 31/7048, 31/7052, A61P 31/04

(21) International Application Number:  
PCT/US2004/036146

(22) International Filing Date: 29 October 2004 (29.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/515,909 30 October 2003 (30.10.2003) US

(71) Applicant (for all designated States except US): **RIB-X  
PHARMACEUTICALS, INC.** [US/US]; 300 George  
Street, Suite 301, New Haven, CT 06511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OYELERE, Adeg-  
boyega, K.** [NG/US]; 149 School Street, Hamden, CT  
06518 (US). **FARMER, Jay, J.** [US/US]; Apt. 4, 198  
Edwards Street, New Haven, CT 06511 (US).

(74) Agents: **GREENHALGH, Duncan, A.** et al.; Patent  
Administrator, Goodwin Procter LLP, Exchange Place, 53  
State Street, Boston, MA 02109 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: BIFUNCTIONAL MACROLIDE HETEROCYCLIC COMPOUNDS AND MEHTODS OF MAKING AND USING  
THE SAME

(57) Abstract: The present invention relates generally to the field of anti-infective, anti-proliferative, anti-inflammatory, and proki-  
netic agents. More particularly, the invention relates to a family of bifunctional compounds useful as therapeutic agents. These  
compounds have both a macrolide ring and at least one heterocyclic moiety. The present invention further relates to processes for  
the preparation of such compounds, to intermediates useful in their preparation, to the use of the compounds as therapeutic agents,  
and to pharmaceuticals compositions containing them.

**BIFUNCTIONAL MACROLIDE HETEROCYCLIC COMPOUNDS AND METHODS OF  
MAKING AND USING THE SAME**

**RELATED APPLICATION**

This application claims the benefit of and priority to U.S. Patent Application No. 60/515,909, filed October 30, 2003, the disclosure of which is incorporated by reference herein.

**FIELD OF THE INVENTION**

5           The present invention relates generally to the field of anti-infective, anti-proliferative, anti-inflammatory, and prokinetic agents, and more particularly, the invention relates to a family of bifunctional macrolide heterocyclic compounds useful as such agents.


**BACKGROUND**

10           Since the discovery of penicillin in the 1920s and streptomycin in the 1940s, many new compounds have been discovered or specifically designed for use as antibiotic agents. It was once believed that infectious diseases could be completely controlled or eradicated with the use of such therapeutic agents. However, such beliefs have been challenged by the fact that strains of microorganisms resistant to currently effective therapeutic agents continue to evolve. Almost every antibiotic agent developed for clinical use has encountered problems with the emergence of  
15           resistant bacteria. For example, resistant strains of Gram-positive bacteria such as methicillin-resistant staphylococci, penicillin-resistant streptococci, and vancomycin-resistant enterococci have developed, and can cause serious and often time fatal results for patients infected with such resistant bacteria. Bacteria that are resistant to the macrolide antibiotics have developed. Also, Gram-negative strains of bacteria such as *H. influenzae* and *M. catarrhalis* have been identified.  
20           See, e.g., F.D. Lowry, Antimicrobial resistance: the example of *Staphylococcus aureus*, *J. Clin. Invest.*, Vol. 111, No. 9, pp. 1265-1273 (2003); and Gold, H.S. and Moellering, R.C., Jr., Antimicrobial-drug resistance. *N. Engl. J. Med.*, vol. 335, 1445-53 (1996).

          This problem of resistance is not limited to the area of anti-infective agents, because resistance has also been encountered with anti-proliferative agents used in cancer chemotherapy.  
25           Therefore, the need exists to develop new anti-infective and anti-proliferative agents that are both effective against resistant bacteria and strains of cells and against which bacteria and strains of cells are less likely to develop resistance.

- 2 -

Despite this problem of increasing antibiotic resistance, no new major classes of antibiotics have been developed for clinical use since the approval in the United States in 2000 of the oxazolidinone ring-containing antibiotic, N-[[[(5S)-3-[3-fluoro-4-(4-morpholinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl acetamide (see structure 1), which is known as linezolid and which is sold under the tradename Zyvox® (see compound A). See, R.C. Moellering, Jr., Linezolid: The First Oxazolidinone Antimicrobial, *Annals of Internal Medicine*, Vol. 138, No. 2, pp. 135-142 (2003).



1

Linezolid was approved for use as an anti-bacterial agent active against Gram-positive organisms. However, linezolid-resistant strains of organisms are already being reported. See Tsiodras *et al.*, *Lancet*, 2001, 358, 207; Gonzales *et al.*, *Lancet*, 2001, 357, 1179; Zurenko *et al.*, *Proceedings Of The 39<sup>th</sup> Annual Interscience Conference On Antibacterial Agents And Chemotherapy (ICAAC)*; San Francisco, CA, USA, September 26-29, 1999). However, investigators have been working to develop other effective linezolid derivatives. Research has indicated that the oxazolidinone ring could be important for linezolid's activity. The literature describes molecules having small groups substituted at the C-5 of the oxazolidinone ring, and early structure-activity relationships suggested that compounds with larger groups at the C-5 position were less active as anti-bacterial agents. As a consequence, investigators have been reluctant to place large substituents at the C-5 position of oxazolidinone rings in developing new anti-microbial agents.

Another class of antibiotics is the macrolides, which is so named for the 14- to 16-membered ring that is the major structural characteristic of this class of compounds. The first macrolide antibiotic to be developed was erythromycin, which was isolated from a soil sample from the Philippines in 1952. Even though erythromycin has been one of the most widely prescribed antibiotics, it has the disadvantages of relatively low bioavailability, gastrointestinal side effects, and a limited spectrum of activity. See Yong-Ji Wu, Highlights of Semi-synthetic Developments from Erythromycin A, *Current Pharm. Design* 6, pp. 181-223 (2000), and Yong-

- 3 -

Ji Wu and Wei-uo Su, Recent Developments on Ketolides and Macrolides, *Curr. Med. Chem.*, 8(14), pp. 1727-1758 (2001).

In the search for new therapeutic agents, pharmaceutical researchers have tried combining or linking various portions of antibiotic molecules. However, this approach has met with limited success.

U.S. Patent No. 5,693,791, to Truett, issued December 2, 1997 describes an antibiotic of the formula:

A-L-B


wherein A and B are antibiotics selected from the group consisting of sulfonamides, penicillins, cephalosporins, quinolones, chloramphenicol, erythromycin (i.e., a macrolide antibiotic), metronidazole, tetracyclines, and aminoglycosides. L is a linker formed from a difunctional linking agent.

PCT publication No. WO 99/63937, to Advanced Medicine, Inc., published December 16, 1999, describes multi-binding compounds useful as antibiotics that are of the following formula:

$(L)_p(X)_q$

wherein L is selected from the group consisting of a macrolide antibiotic, an aminoglycoside, lincosamide, oxazolidinone, streptogramin, tetracycline, or another compound that binds to bacterial ribosomal RNA and/or to one or more proteins involved in ribosomal protein synthesis in the bacterium. In the formula, p is an integer from 2-10, q is an integer from 1-20, and X is a linker.

U.S. Patent No. 6,034,069, to Or et al., issued March 7, 2000 depicts a series of 3'-N-modified 6-O-substituted erythromycin ketolide derivatives such as shown below. R, R<sup>1</sup>, and R<sup>2</sup> are selected from the group consisting of a variety of groups, including aryl-alkoxy-heteroaryl-alkylene. R<sup>p</sup> is H or a hydroxy protecting group. W is absent or is O, NH, or NCH<sub>3</sub>. R<sup>w</sup> is H or an optionally substituted alkyl group.



- 4 -


Notwithstanding the foregoing, there is an ongoing need for new anti-infective and anti-proliferative agents. In the present invention it has been found that new macrolide heterocyclic agents, including agents with relatively large substituents at the C5 position of an oxazolidinone ring and at similar positions in other heterocyclic rings can be prepared having good activity.

Furthermore, because many anti-infective and anti-proliferative agents have utility as anti-inflammatory agents and also as prokinetic (gastrointestinal modulatory) agents, there is also an ongoing need for new compounds useful as anti-inflammatory and prokinetic agents. The present invention provides compounds which meet these needs.


### SUMMARY OF THE INVENTION

The present invention provides compounds useful as anti-infective agents and/or anti-proliferative agents, for example anti-microbial agents, anti-acterial agents, anti-iotic agents, anti-fungal agents, anti-parasitic agents, anti-viral agents, and chemotherapeutic agents. The present invention also provides compounds useful as anti-inflammatory agents, and/or prokinetic (i.e. gastrointestinal modulatory) agents. The present invention also provides pharmaceutically acceptable salts, ester, or prodrugs thereof.

The present invention provides compounds having both a macrolide ring and at least one heterocyclic moiety having the formula:



or a stereoisomer, or pharmaceutically acceptable salt, ester or prodrug thereof, wherein D-Het is selected from the group consisting of:



B is selected from the group consisting of:

-C(O)NH-, -C(S)NH-, -NHC(O)-, -NHC(S)-, -S(O)<sub>2</sub>NH-, -NHS(O)<sub>2</sub>-,  
 -OC(O)NH-, -OC(S)NH-, -NHC(O)NH-, -NHC(S)NH-, -NHC(O)O-,  
 -NHC(S)O-, and -NR<sup>11</sup>-;

n is 0 or 1, and the variables A, D, E, M, R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6'</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, and R<sup>10</sup> can be selected from the group consisting of the respective chemical moieties later defined in the detailed description.

In addition, the invention provides methods of synthesizing the foregoing compounds and useful chemical intermediates for synthesizing the foregoing compounds. Following synthesis, a therapeutically effective amount of one or more of the compounds can be formulated with a pharmaceutically acceptable carrier for administration to a mammal for use as an anti-cancer, anti-microbial, anti-biotic, anti-fungal, anti-parasitic or anti-viral agent, or to treat a proliferative disease, an inflammatory disease or a gastrointestinal motility disorder. Accordingly, the compounds or the formulations can be administered, for example, via oral, parenteral, or topical routes, to provide an effective amount of the compound to the mammal.

These and other aspects and embodiments of the invention can be more fully understood by reference to the following detailed description and claims.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a family of compounds that can be used as anti-proliferative agents and/or anti-infective agents. The compounds may be used without limitation, for example, as anti-cancer, anti-microbial, anti-bacterial, anti-fungal, anti-parasitic and/or anti-viral agents. Further, the present invention provides a family of compounds that can be used without limitation as anti-inflammatory agents, for example, for use in treating chronic inflammatory airway diseases, and/or as prokinetic agents, for example, for use in treating gastrointestinal motility disorders such as gastroesophageal reflux disease, gastroparesis (diabetic and post surgical), irritable bowel syndrome, and constipation.

The compounds described herein may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic, and geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated. All processes used to prepare compounds of the present invention and intermediates made therein are considered to be part of the present invention. All tautomers of shown or described compounds are also considered to be part of the present invention.

### 1. Definitions

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (e.g., C=C, C=N, or N=N). The present invention, in general, does not cover groups such as N-halo, S(O)H, and SO<sub>2</sub>H.

The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass

numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

When any variable (e.g.,  $R^3$ ) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other  
5 occurrence. Thus, for example, if a group is shown to be substituted with 0-2  $R^3$  moieties, then the group may optionally be substituted with up to two  $R^3$  moieties and  $R^3$  at each occurrence is selected independently from the definition of  $R^3$ . Also, combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring,  
10 then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

15 In cases wherein there are nitrogens in the compounds of the present invention, these can be converted to N-oxides by treatment with an oxidizing agent (e.g., MCPBA and/or hydrogen peroxides) to afford other compounds of the present invention. Thus, all shown and claimed nitrogens are considered to cover both the shown nitrogen and its N-oxide ( $N \rightarrow O$ ) derivative.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated  
20 aliphatic hydrocarbon groups having the specified number of carbon atoms.  $C_{1-6}$  alkyl is intended to include  $C_1$ ,  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$ , and  $C_6$  alkyl groups.  $C_{1-8}$  alkyl is intended to include  $C_1$ ,  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$ ,  $C_6$ ,  $C_7$ , and  $C_8$  alkyl groups. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl, n-hexyl, n-heptyl, and n-octyl.

25 As used herein, "alkenyl" is intended to include hydrocarbon chains of either straight or branched configuration and one or more unsaturated carbon-carbon bonds that may occur in any stable point along the chain, such as ethenyl and propenyl.  $C_{2-6}$  alkenyl is intended to include  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$ , and  $C_6$  alkenyl groups.  $C_{2-8}$  alkenyl is intended to include  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$ ,  $C_6$ ,  $C_7$ , and  $C_8$  alkenyl groups.

30 As used herein, "alkynyl" is intended to include hydrocarbon chains of either straight or branched configuration and one or more triple carbon-carbon bonds that may occur in any stable point along the chain, such as ethynyl and propynyl.  $C_{2-6}$  alkynyl is intended to include  $C_2$ ,  $C_3$ ,



C<sub>4</sub>, C<sub>5</sub>, and C<sub>6</sub> alkynyl groups. C<sub>2-8</sub> alkynyl is intended to include C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, C<sub>6</sub>, C<sub>7</sub>, and C<sub>8</sub> alkynyl groups.

As used herein, "cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, or cyclopentyl. C<sub>3-8</sub> cycloalkyl is intended to include C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, C<sub>6</sub>,  
5 C<sub>7</sub>, and C<sub>8</sub> cycloalkyl groups.

As used herein, "halo" or "halogen" refers to fluoro, chloro, bromo, and iodo. "Counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, and sulfate.

As used herein, "haloalkyl" is intended to include both branched and straight-chain  
10 saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example -C<sub>v</sub>F<sub>w</sub> where v = 1 to 3 and w = 1 to (2v+1)). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl.

As used herein, "alkoxy" refers to an alkyl group as defined above with the indicated  
15 number of carbon atoms attached through an oxygen bridge. C<sub>1-6</sub> alkoxy, is intended to include C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, and C<sub>6</sub> alkoxy groups. C<sub>1-8</sub> alkoxy, is intended to include C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, C<sub>6</sub>, C<sub>7</sub>, and C<sub>8</sub> alkoxy groups. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, s-pentoxy, n-heptoxy, and n-octoxy.

As used herein, "alkylthio" refers to an alkyl group as defined above with the indicated  
20 number of carbon atoms attached through a sulfur bridge. C<sub>1-6</sub> alkylthio, is intended to include C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, and C<sub>6</sub> alkylthio groups. C<sub>1-8</sub> alkylthio, is intended to include C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub>, C<sub>4</sub>, C<sub>5</sub>, C<sub>6</sub>, C<sub>7</sub>, and C<sub>8</sub> alkylthio groups.

As used herein, "carbocycle" or "carbocyclic ring" is intended to mean, unless otherwise  
25 specified, any stable 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12-membered monocyclic, bicyclic or tricyclic ring, any of which may be saturated, unsaturated, or aromatic, recognizing that rings with certain numbers of members cannot be bicyclic or tricyclic, e.g., a 3-membered ring can only be a monocyclic ring. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, [3.3.0]bicyclooctane,  
30 [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl. As shown above, bridged rings are also included in

the definition of carbocycle (e.g., [2.2.2]bicyclooctane). A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge. Fused (e.g., naphthyl and tetrahydronaphthyl) and spiro rings are also included.

As used herein, the term "heterocycle" means, unless otherwise stated, a stable 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12-membered monocyclic, bicyclic or tricyclic heterocyclic ring (recognizing that rings with certain numbers of members cannot be bicyclic or tricyclic, e.g., a 3-membered ring can only be a monocyclic ring), which is saturated, unsaturated, or aromatic, and consists of carbon atoms and one or more ring heteroatoms, e.g., 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen, and sulfur, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a second ring (e.g., a benzene ring). The nitrogen and sulfur heteroatoms may optionally be oxidized (i.e.,  $N \rightarrow O$  and  $S(O)_p$ , where  $p = 1$  or  $2$ ). When a nitrogen atom is included in the ring it is either N or NH, depending on whether or not it is attached to a double bond in the ring (i.e., a hydrogen is present if needed to maintain the tri-valency of the nitrogen atom). The nitrogen atom may be substituted or unsubstituted (i.e., N or NR wherein R is H or another substituent, as defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more atoms (i.e., C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Preferred bridges include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms, and a carbon-nitrogen group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge. Spiro and fused rings are also included.

As used herein, the term "aromatic heterocycle" or "heteroaryl" is intended to mean a stable 5, 6, 7, 8, 9, 10, 11, or 12-membered monocyclic or bicyclic heterocyclic aromatic ring (recognizing that rings with certain numbers of members cannot be a bicyclic aromatic, e.g., a 5-membered ring can only be a monocyclic aromatic ring), which consists of carbon atoms and one

or more heteroatoms, e.g., 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen, and sulfur. In the case of bicyclic heterocyclic aromatic rings, only one of the two rings needs to be aromatic (e.g., 2,3-dihydroindole), though both may be (e.g., quinoline). The second ring can also be fused or  
5 bridged as defined above for heterocycles. The nitrogen atom may be substituted or unsubstituted (i.e., N or NR wherein R is H or another substituent, as defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (i.e., N→O and S(O)<sub>p</sub>, where p = 1 or 2). It is to be noted that total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, acridinyl, azocinyl,  
10 benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazoliny, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazoliny, carbazolyl, 4*aH*-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2*H*,6*H*-1,5,2-dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, 1*H*-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3*H*-indolyl,  
15 isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenaziny, phenothiazinyl, phenoxathinyl, phenoxazinyl, phthalazinyl,  
20 piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridoazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2*H*-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4*H*-quinoliziny, quinoxalinyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, tetrazolyl, 6*H*-1,2,5-thiadiazinyl,  
25 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

As used herein, the phrase "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical  
30 judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 2-acetoxybenzoic, 2-hydroxyethane sulfonic, acetic, ascorbic, benzene sulfonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulfonic, ethane sulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycollyarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodide, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulfonic, maleic, malic, mandelic, methane sulfonic, napsylic, nitric, oxalic, pamoic, pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicyclic, stearic, subacetic, succinic, sulfamic, sulfanilic, sulfuric, tannic, tartaric, and toluene sulfonic.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing Company, Easton, PA, 1990, 1445.

Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.) the compounds of the present invention may be delivered in prodrug form. Thus, the present invention is intended to cover prodrugs of the presently claimed compounds, methods of delivering the same and compositions containing the same. "Prodrugs" are intended to include any covalently bonded carriers that release an active parent drug of the present invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug of the present invention is administered to a mammalian subject, it cleaves to form a free hydroxyl, free amino,

or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate, and benzoate derivatives of alcohol and amine functional groups in the compounds of the present invention.

"Stable compound" and "stable structure" are meant to indicate a compound that is  
5 sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. It is preferred that the presently recited compounds do not contain a N-halo, S(O)<sub>2</sub>H, or S(O)H group.

As used herein, "treating" or "treatment" means the treatment of a disease-state in a mammal, particularly in a human, and include: (a) preventing the disease-state from occurring in  
10 a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, i.e., arresting its development; and/or (c) relieving the disease-state, i.e., causing regression of the disease state.

As used herein, "mammal" refers to human and non-human patients.


As used herein, the term "therapeutically effective amount" refers to an amount of a  
15 compound, or a combination of compounds, of the present invention effective when administered alone or in combination as an anti-proliferative and/or anti-infective agent. The combination of compounds is preferably a synergistic combination. Synergy, as described, for example, by Chou and Talalay, *Adv. Enzyme Regul.* 1984, 22:27-55, occurs when the effect of the compounds when administered in combination is greater than the additive effect of the compounds when  
20 administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at sub-optimal concentrations of the compounds. Synergy can be in terms of lower cytotoxicity, increased anti-proliferative and/or anti-infective effect, or some other beneficial effect of the combination compared with the individual components.

All percentages and ratios used herein, unless otherwise indicated, are by weight.

25 Throughout the description, where compositions are described as having, including, or comprising specific components, or where processes are described as having, including, or comprising specific process steps, it is contemplated that compositions of the present invention also consist essentially of, or consist of, the recited components, and that the processes of the present invention also consist essentially of, or consist of, the recited processing steps. Further, it  
30 should be understood that the order of steps or order for performing certain actions are immaterial so long as the invention remains operable. Moreover, two or more steps or actions may be conducted simultaneously.

## 2. Compounds of the Invention


In one aspect, the invention provides compounds having the formula:



5 or pharmaceutically acceptable salt, ester or prodrug thereof,

wherein

D-Het is selected from the group consisting of:



A is selected from the group consisting of:

a) carbonyl, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl d) -C(O)-C<sub>1-6</sub> alkyl, and  
e) -C(O)-C<sub>2-6</sub> alkenyl,

wherein

- 5 i) 0-2 carbon atoms of the C<sub>1-6</sub> alkyl and C<sub>2-6</sub> alkenyl groups in any of b) – e) optionally are replaced by a moiety selected from the group consisting of O, S(O)<sub>p</sub>, and NR<sup>11</sup>, and  
ii) any of b) – e) optionally is substituted with one or more R<sup>12</sup> groups;

B is selected from the group consisting of:

- 10 a) -C(O)NH-, b) -C(S)NH-, c) -NHC(O)-, d) -NHC(S)-, e) -S(O)<sub>2</sub>NH-,  
f) -NHS(O)<sub>2</sub>-, g) -OC(O)NH-, h) -OC(S)NH-, i) -NHC(O)NH-, j) -NHC(S)NH-,  
k) -NHC(O)O-, l) -NHC(S)O-, and m) -NR<sup>11</sup>-;


n is 0 or 1;

D is selected from the group consisting of:


- 15 a) -CH<sub>2</sub>-, b) -C(O)-, c) -C(S)-, d) -C(=NOR<sup>11</sup>)-, e) -CH<sub>2</sub>CH<sub>2</sub>-, f) -OCH<sub>2</sub>-,  
g) -SCH<sub>2</sub>-, h) -S(O)CH<sub>2</sub>-, i) -S(O)<sub>2</sub>CH<sub>2</sub>-, j) -NR<sup>11</sup>CH<sub>2</sub>-, k) -C(O)CH<sub>2</sub>-,  
l) -C(S)CH<sub>2</sub>-, and m) -C(=NOR<sup>11</sup>)CH<sub>2</sub>-;

E is selected from the group consisting of:


a)



b)



c)



d) 5-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally substituted with one or more  $R^{12}$  groups;

e)  $C_{5-10}$  saturated, unsaturated, or aromatic carbocycle, optionally substituted with one or more  $R^{12}$  groups;

f)  $C_{1-8}$  alkyl,

g)  $C_{2-8}$  alkenyl,

h)  $C_{2-8}$  alkynyl,

i)  $C_{1-8}$  alkoxy,

j)  $C_{1-8}$  alkylthio,

k)  $C_{1-8}$  acyl,

l)  $S(O)_r R^{11}$ ; and

m) hydrogen,

wherein any of f) – k) optionally is substituted with

i) one or more  $R^{12}$  groups;

ii) 5-6 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally substituted with one or more  $R^{12}$  groups; or

iii)  $C_{5-10}$  saturated, unsaturated, or aromatic carbocycle, optionally substituted with one or more  $R^{12}$  groups;

M is selected from the group consisting of:

a)  $-C(O)-$ , b)  $-C(=NOR^{11})-$ , c)  $-CH(-OR^{11})-$ , d)  $-NR^{11}-CH_2-$ , e)  $-CH_2-NR^{11}-$ ,

f)  $-CH(NR^{11}R^{11})-$ , g)  $-C(=NNR^{11}R^{11})-$ , h)  $-NR^{11}-C(O)-$ , i)  $-C(O)NR^{11}-$ , and

j)  $-C(=NR^{11})-$ ;

R is selected from the group consisting of H and  $C_{1-6}$  alkyl;

$R^1$  is selected from the group consisting of:

a) H, b) Cl, c) F, d) Br, e) I, f)  $-NR^{11}R^{11}$  g)  $-NR^{11}C(O)R^{11}$ , h)  $-OR^{11}$ ,

i)  $-OC(O)R^{11}$ , j)  $-OC(O)OR^{11}$ , k)  $-OC(O)NR^{11}R^{11}$ , l)  $-O-C_{1-6}$  alkyl- $R^{12}$ ,




m)  $-\text{OC}(\text{O})-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , n)  $-\text{OC}(\text{O})\text{O}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ ,  
 o)  $-\text{OC}(\text{O})\text{NR}^{11}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , p)  $\text{C}_{1-6}$  alkyl, q)  $\text{C}_{1-6}$  alkenyl, r)  $\text{C}_{1-6}$  alkynyl,  
 wherein any of l) – r) optionally is substituted with one or more  $\text{R}^{12}$   
 groups;

5  $\text{R}^2$  is H;

$\text{R}^3$  is selected from the group consisting of:

a) H, b)  $-\text{OR}^{11}$ , c)  $-\text{O}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , d)  $-\text{OC}(\text{O})\text{R}^{11}$ , e)  $-\text{OC}(\text{O})-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ ,  
 f)  $-\text{OC}(\text{O})\text{OR}^{11}$ , g)  $-\text{OC}(\text{O})\text{O}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , h)  $-\text{OC}(\text{O})\text{NR}^{11}\text{R}^{11}$ ,  
 i)  $-\text{OC}(\text{O})\text{NR}^{11}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , and  
 10 j)




alternatively,  $\text{R}^2$  and  $\text{R}^3$  taken together form a carbonyl group;

$\text{R}^4$  is selected from the group consisting of:

a) H, b)  $\text{R}^{11}$ , c)  $-\text{C}(\text{O})\text{R}^{11}$  d)  $-\text{C}(\text{O})\text{OR}^{11}$  e)  $-\text{C}(\text{O})\text{NR}^{11}\text{R}^{11}$ , f)  $-\text{C}_{1-6}$  alkyl- $\text{G}-\text{R}^{11}$ ,  
 15 g)  $-\text{C}_{2-6}$  alkenyl- $\text{G}-\text{R}^{11}$ , and h)  $-\text{C}_{2-6}$  alkynyl- $\text{G}-\text{R}^{11}$ ;

alternatively  $\text{R}^3$  and  $\text{R}^4$ , taken together with the atoms to which they are bonded, form:



G is selected from the group consisting of:


a)  $-\text{C}(\text{O})-$ , b)  $-\text{C}(\text{O})\text{O}-$ , c)  $-\text{C}(\text{O})\text{NR}^{11}-$ , d)  $-\text{C}(=\text{NR}^{11})-$ , e)  $-\text{C}(=\text{NR}^{11})\text{O}-$ ,  
 20 f)  $-\text{C}(=\text{NR}^{11})\text{NR}^{11}-$ , g)  $-\text{OC}(\text{O})-$ , h)  $-\text{OC}(\text{O})\text{O}-$ , i)  $-\text{OC}(\text{O})\text{NR}^{11}-$ , j)  $-\text{NR}^{11}\text{C}(\text{O})-$ ,  
 k)  $-\text{NR}^{11}\text{C}(\text{O})\text{O}-$ , l)  $-\text{NR}^{11}\text{C}(\text{O})\text{NR}^{11}-$ , m)  $-\text{NR}^{11}\text{C}(=\text{NR}^{11})\text{NR}^{11}-$ , and o)  $-\text{S}(\text{O})_p-$ ;

$\text{R}^5$  is selected from the group consisting of:

a)  $\text{R}^{11}$ , b)  $-\text{OR}^{11}$ , c)  $-\text{NR}^{11}\text{R}^{11}$ , d)  $-\text{O}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , e)  $-\text{C}(\text{O})-\text{R}^{11}$ ,  
 f)  $-\text{C}(\text{O})-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , g)  $-\text{OC}(\text{O})-\text{R}^{11}$ , h)  $-\text{OC}(\text{O})-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ ,  
 25 i)  $-\text{OC}(\text{O})\text{O}-\text{R}^{11}$ , j)  $-\text{OC}(\text{O})\text{O}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , k)  $-\text{OC}(\text{O})\text{NR}^{11}\text{R}^{11}$ ,

- l)  $-\text{OC}(\text{O})\text{NR}^{11}-\text{C}_{1-6}$  alkyl- $\text{R}^{12}$ , m)  $-\text{C}(\text{O})-\text{C}_{2-6}$  alkenyl- $\text{R}^{12}$ , and  
 n)  $-\text{C}(\text{O})-\text{C}_{2-6}$  alkynyl- $\text{R}^{12}$ ;

alternatively,  $\text{R}^4$  and  $\text{R}^5$ , taken together with the atoms to which they are bonded, form:



5 wherein

Q is CH or N, and

$\text{R}^{23}$  is  $-\text{OR}^{11}$ , or  $\text{R}^{11}$ ;


$\text{R}^6$  is selected from the group consisting of:

- a)  $-\text{OR}^{11}$ , b)  $-\text{C}_{1-6}$  alkoxy- $\text{R}^{12}$ , c)  $-\text{C}(\text{O})\text{R}^{11}$ , d)  $-\text{OC}(\text{O})\text{R}^{11}$ , e)  $-\text{OC}(\text{O})\text{OR}^{11}$ ,  
 10 f)  $-\text{OC}(\text{O})\text{NR}^{11}\text{R}^{11}$ , and g)  $-\text{NR}^{11}\text{R}^{11}$ ;

alternatively,  $\text{R}^5$  and  $\text{R}^6$  taken together with the atoms to which they are attached form a  
 5-membered ring by attachment to each other through a linker selected from the group consisting  
 of:

- a)  $-\text{OC}(\text{R}^{12})_2\text{O}-$ , b)  $-\text{OC}(\text{O})\text{O}-$ , c)  $-\text{OC}(\text{O})\text{NR}^{11}-$ , d)  $-\text{NR}^{11}\text{C}(\text{O})\text{O}-$ ,  
 15 e)  $-\text{OC}(\text{O})\text{NOR}^{11}-$ , f)  $-\text{NOR}^{11}-\text{C}(\text{O})\text{O}-$ , g)  $-\text{OC}(\text{O})\text{NNR}^{11}\text{R}^{11}-$ ,  
 h)  $-\text{NNR}^{11}\text{R}^{11}-\text{C}(\text{O})\text{O}-$ , i)  $-\text{OC}(\text{O})\text{C}(\text{R}^{12})_2-$ , j)  $-\text{C}(\text{R}^{12})_2\text{C}(\text{O})\text{O}-$ , k)  $-\text{OC}(\text{S})\text{O}-$ ,  
 l)  $-\text{OC}(\text{S})\text{NR}^{11}-$ , m)  $-\text{NR}^{11}\text{C}(\text{S})\text{O}-$ , n)  $-\text{OC}(\text{S})\text{NOR}^{11}-$ , o)  $-\text{NOR}^{11}-\text{C}(\text{S})\text{O}-$ ,  
 p)  $-\text{OC}(\text{S})\text{NNR}^{11}\text{R}^{11}-$ , q)  $-\text{NNR}^{11}\text{R}^{11}-\text{C}(\text{S})\text{O}-$ , r)  $-\text{OC}(\text{S})\text{C}(\text{R}^{12})_2-$ , and  
 s)  $-\text{C}(\text{R}^{12})_2\text{C}(\text{S})\text{O}-$ ;

20 alternatively, M,  $\text{R}^5$ , and  $\text{R}^6$  taken together with the atoms to which they are attached  
 form:



wherein J is selected from the group consisting of O and  $\text{NR}^{11}$ ;

$\text{R}^{6'}$  is selected from the group consisting of

5 a) -H, b) -C<sub>1-4</sub> alkyl, c) C<sub>2-4</sub> alkenyl, which can be further substituted with C<sub>1-12</sub> alkyl or one or more halogens, d) C<sub>2-4</sub> alkynyl, which can be further substituted with C<sub>1-12</sub> alkyl or one or more halogens, e) aryl or heteroaryl, which can be further substituted with C<sub>1-12</sub> alkyl or one or more halogens, f) -C(O)H, g) -COOH, h) -CN, i) -COOR<sup>11</sup>, j) -C(O)NR<sup>11</sup>R<sup>11</sup>, k) -C(O)R<sup>11</sup>, and l) -C(O)SR<sup>11</sup>,  
 10 wherein b) is further substituted with one or more substituents selected from the group consisting of aa) -OR<sup>11</sup>, bb) halogen, cc) -SR<sup>11</sup>, dd) C<sub>1-12</sub> alkyl, which can be further substituted with halogen, hydroxyl, C<sub>1-6</sub> alkoxy, or amino, ee) -OR<sup>11</sup>, ff) -SR<sup>11</sup>, gg) -NR<sup>11</sup>R<sup>11</sup>, hh) -CN, ii) -NO<sub>2</sub>, jj) -NC(O)R<sup>11</sup>, kk) -COOR<sup>11</sup>, ll) -N<sub>3</sub>, mm) =N-O-R<sup>11</sup>, nn) =NR<sup>11</sup>, oo) =N-NR<sup>11</sup>R<sup>11</sup>, pp) =N-NH-C(O)R<sup>11</sup>, and qq) =N-NH-C(O)NR<sup>11</sup>R<sup>11</sup>;

alternatively R<sub>6</sub> and R<sub>6'</sub> are taken together with the atom to which they are attached to form an epoxide, a carbonyl, an olefin, or a substituted olefin, or a C<sub>3</sub>-C<sub>7</sub> carbocyclic, carbonate, or carbamate, wherein the nitrogen of said carbamate can be further substituted with a C<sub>1</sub>-C<sub>6</sub>  
 15 alkyl;

R<sup>7</sup> is selected from the group consisting of:

a) C<sub>1-6</sub> alkyl, b) C<sub>2-6</sub> alkenyl, and c) C<sub>2-6</sub> alkynyl,

wherein any of a) - c) optionally is substituted with one or more R<sup>12</sup> groups;

20 R<sup>8</sup> is selected from the group consisting of H and -C(O)R<sup>11</sup>;

R<sup>9</sup> is selected from the group consisting of H, OH, and -OR<sup>11</sup>;

R<sup>10</sup> is selected from the group consisting of:

a) H, b) R<sup>11</sup>, c) -C<sub>1-6</sub> alkyl-G-R<sup>12</sup>, d) -C<sub>2-6</sub> alkenyl-G-R<sup>12</sup>, and

e) -C<sub>2-6</sub> alkynyl-G-R<sup>12</sup>,

25 wherein the C<sub>1-6</sub>-alkyl, C<sub>2-6</sub> alkenyl, and C<sub>2-6</sub> alkynyl group in any of c) - e) optionally is substituted with one or more R<sup>12</sup> groups;

R<sup>11</sup>, at each occurrence, independently is selected from the group consisting of:

a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>6-10</sub> saturated, unsaturated, or aromatic carbocycle, f) 3-12 membered saturated, unsaturated, or aromatic

30 heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, g) -C(O)-C<sub>1-6</sub> alkyl,

h) -C(O)-C<sub>2-6</sub> alkenyl, i) -C(O)-C<sub>2-6</sub> alkynyl, j) -C(O)-C<sub>6-10</sub> saturated, unsaturated,

- or aromatic carbocycle, k) -C(O)-3-12 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, l) -C(O)O-C<sub>1-6</sub> alkyl, m) -C(O)O-C<sub>2-6</sub> alkenyl, n) -C(O)O-C<sub>2-6</sub> alkynyl, o) -C(O)O-C<sub>6-10</sub> saturated, 5 unsaturated, or aromatic carbocycle, p) -C(O)O-3-12 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and q) -C(O)NR<sup>13</sup>R<sup>13</sup>, wherein any of b) – p) optionally is substituted with one or more R<sup>12</sup> groups,
- 10 alternatively, NR<sup>11</sup>R<sup>11</sup> forms a 3-7 membered saturated, unsaturated or aromatic ring including the nitrogen atom to which the R<sup>11</sup> groups are bonded and optionally one or more moieties selected from the group consisting of: O, S(O)<sub>p</sub>, and NR<sup>15</sup>;
- R<sup>12</sup> is selected from the group consisting of:
- 15 a) R<sup>14</sup>, b) C<sub>1-8</sub> alkyl, c) C<sub>2-8</sub> alkenyl, d) C<sub>2-8</sub> alkynyl, e) C<sub>3-12</sub> saturated, unsaturated, or aromatic carbocycle, f) 3-12 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and g) -NR<sup>15</sup>C(O)OR<sup>15</sup>, wherein any of b) – f) optionally is substituted with one or more R<sup>14</sup> groups;
- 20 R<sup>13</sup>, at each occurrence, independently is selected from the group consisting of:
- a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>3-10</sub> saturated, unsaturated, or aromatic carbocycle, and f) 3-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,
- 25 wherein any of b) – f) optionally is substituted with one or more moieties selected from the group consisting of:
- 30 carbonyl; formyl; F; Cl; Br; I; CN; NO<sub>2</sub>; OR<sup>15</sup>; -S(O)<sub>p</sub>R<sup>15</sup>; -C(O)R<sup>15</sup>; -C(O)OR<sup>15</sup>; -OC(O)R<sup>15</sup>; -C(O)NR<sup>15</sup>R<sup>15</sup>; -OC(O)NR<sup>15</sup>R<sup>15</sup>; -C(=NR<sup>15</sup>)R<sup>15</sup>; -C(R<sup>15</sup>)(R<sup>15</sup>)OR<sup>15</sup>; -C(R<sup>15</sup>)<sub>2</sub>OC(O)R<sup>15</sup>; -C(R<sup>15</sup>)(OR<sup>15</sup>)(CH<sub>2</sub>)<sub>r</sub>NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>OR<sup>15</sup>; -NR<sup>15</sup>C(O)R<sup>15</sup>; -NR<sup>15</sup>C(O)OR<sup>15</sup>; -NR<sup>15</sup>C(O)NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>S(O)<sub>r</sub>R<sup>15</sup>; -C(OR<sup>15</sup>)(OR<sup>15</sup>)R<sup>15</sup>; -C(R<sup>15</sup>)<sub>2</sub>NR<sup>15</sup>R<sup>15</sup>; =NR<sup>15</sup>; -C(S)NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>C(S)R<sup>15</sup>; -OC(S)NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>C(S)OR<sup>15</sup>;

5  
 -NR<sup>15</sup>C(S)NR<sup>15</sup>R<sup>15</sup>; -SC(O)R<sup>15</sup>; C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl;  
 C<sub>2-8</sub> alkynyl; C<sub>1-8</sub> alkoxy; C<sub>1-8</sub> alkylthio; C<sub>1-8</sub> acyl; saturated,  
 unsaturated, or aromatic C<sub>3-10</sub> carbocycle; and saturated,  
 unsaturated, or aromatic 3-10 membered heterocycle containing  
 one or more heteroatoms selected from the group consisting of  
 nitrogen, oxygen, and sulfur,

alternatively, NR<sup>13</sup>R<sup>13</sup> forms a 3-10 membered saturated, unsaturated or aromatic ring  
 including the nitrogen atom to which the R<sup>13</sup> groups are attached and optionally one or more  
 moieties selected from the group consisting of O, S(O)<sub>p</sub>, NR<sup>15</sup>, and N;

10 alternatively, CR<sup>13</sup>R<sup>13</sup> forms a carbonyl group;

R<sup>14</sup>, at each occurrence, is selected from the group consisting of:

- a) H, b) carbonyl, c) F, d) Cl, e) Br, f) I, g) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CF<sub>3</sub>, h) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CN,  
 i) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NO<sub>2</sub>, j) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, k) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OR<sup>16</sup>,  
 l) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>S(O)<sub>p</sub>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, m) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 15 n) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OC(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, o) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>SC(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 p) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)O(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, q) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>C(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 r) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, s) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NR<sup>13</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 t) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NNR<sup>13</sup>R<sup>13</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 u) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NNR<sup>13</sup>C(O)R<sup>13</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 20 v) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NOR<sup>16</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 w) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>C(O)O(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 x) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OC(O)NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 y) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>C(O)NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 z) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>S(O)<sub>p</sub>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 25 aa) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>S(O)<sub>p</sub>NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 bb) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>S(O)<sub>p</sub>NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, cc) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>R<sup>13</sup>,  
 dd) C<sub>1-6</sub> alkyl, ee) C<sub>2-6</sub> alkenyl, ff) C<sub>2-6</sub> alkynyl, gg) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>-C<sub>3-10</sub> saturated,  
 unsaturated, or aromatic carbocycle, and hh) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>-3-10 membered  
 saturated, unsaturated, or aromatic heterocycle containing one or more  
 30 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,

wherein any of dd) – hh) optionally is substituted with one or more R<sup>16</sup> groups;

alternatively, two R<sup>14</sup> groups may form -O(CH<sub>2</sub>)<sub>s</sub>O-;

R<sup>15</sup> is selected from the group consisting of:

- 5 a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>3-10</sub> saturated, unsaturated, or aromatic carbocycle, f) 3-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, g) -C(O)-C<sub>1-6</sub> alkyl,
- 10 h) -C(O)-C<sub>1-6</sub> alkenyl, g) -C(O)-C<sub>1-6</sub> alkynyl, i) -C(O)-C<sub>3-10</sub> saturated, unsaturated, or aromatic carbocycle, and j) -C(O)-3-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,
- wherein any of b) – j) optionally is substituted with one or more moieties selected from the group consisting of H; F; Cl; Br; I; CN; NO<sub>2</sub>; OH; NH<sub>2</sub>;
- 15 NH(C<sub>1-6</sub> alkyl); N(C<sub>1-6</sub> alkyl)<sub>2</sub>; C<sub>1-6</sub> alkoxy; aryl; substituted aryl; heteroaryl; substituted heteroaryl; and C<sub>1-6</sub> alkyl, optionally substituted with one or more moieties selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, F, Cl, Br, I, CN, NO<sub>2</sub>, and OH;

20 R<sup>16</sup>, at each occurrence, independently is selected from the group consisting of:

- a) R<sup>17</sup>, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) -C<sub>3-10</sub> saturated, unsaturated, or aromatic carbocycle, and f) -3-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,
- 25 wherein any of b) – f) optionally is substituted with one or more R<sup>17</sup> groups;

R<sup>17</sup>, at each occurrence, independently is selected from the group consisting of:

- a) H, b) carbonyl, c) F, d) Cl, e) Br, f) I, g) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CF<sub>3</sub>, h) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CN,
- i) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NO<sub>2</sub>, j) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>R<sup>13</sup>, k) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OR<sup>11</sup>,
- 30 l) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>S(O)<sub>p</sub>R<sup>13</sup>, m) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)R<sup>13</sup>, n) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)OR<sup>13</sup>, o) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OC(O)R<sup>13</sup>, p) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>C(O)R<sup>13</sup>,
- q) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)NR<sup>13</sup>R<sup>13</sup>, r) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NR<sup>13</sup>)R<sup>13</sup>,

s)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{C}(\text{O})\text{NR}^{13}\text{R}^{13}$ , t)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{S}(\text{O})_p\text{R}^{13}$ ,  
 u)  $(\text{CR}^{13}\text{R}^{13})_r\text{S}(\text{O})_p\text{NR}^{13}\text{R}^{13}$ , v)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{S}(\text{O})_p\text{NR}^{13}\text{R}^{13}$ , w)  $\text{C}_{1-6}$  alkyl,  
 x)  $\text{C}_{2-6}$  alkenyl, y)  $\text{C}_{2-6}$  alkynyl, z)  $(\text{CR}^{13}\text{R}^{13})_r\text{-C}_{3-10}$  saturated, unsaturated, or  
 aromatic carbocycle, and aa)  $(\text{CR}^{13}\text{R}^{13})_r\text{-3-10}$  membered saturated, unsaturated,  
 or aromatic heterocycle containing one or more heteroatoms selected from the  
 group consisting of nitrogen, oxygen, and sulfur,

wherein any of w) – aa) optionally is substituted with one or more  
 moieties selected from the group consisting of  $\text{R}^{13}$ ; F; Cl; Br; I; CN;  $\text{NO}_2$ ;  
 $-\text{OR}^{13}$ ;  $-\text{NH}_2$ ;  $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$ ;  $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$ ;  $\text{C}_{1-6}$  alkoxy;  $\text{C}_{1-6}$  alkylthio;  
 and  $\text{C}_{1-6}$  acyl;

$\text{R}^{18}$ , at each occurrence, independently is selected from the group consisting of:

a) H, b)  $\text{OR}^{15}$ , c)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{OC}(\text{O})\text{R}^{15}$ , d)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{OC}(\text{O})\text{OR}^{15}$ ,  
 e)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{OC}(\text{O})\text{NR}^{15}\text{R}^{15}$ , f)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ ,  
 g)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{NR}^{15}\text{C}(\text{O})\text{R}^{15}$ , h)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{NR}^{15}\text{C}(\text{O})\text{OR}^{15}$ ,  
 i)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{NR}^{15}\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ , j)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{NR}^{15}\text{C}(=\text{NH})\text{NR}^{15}\text{R}^{15}$ ,  
 k)  $-\text{O}-\text{C}_{1-6} \text{ alkyl}-\text{S}(\text{O})_p\text{R}^{15}$ , l)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{OC}(\text{O})\text{R}^{15}$ ,  
 m)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{OC}(\text{O})\text{OR}^{15}$ , n)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{OC}(\text{O})\text{NR}^{15}\text{R}^{15}$ ,  
 o)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ , p)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{NR}^{15}\text{C}(\text{O})\text{R}^{15}$ ,  
 q)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{NR}^{15}\text{C}(\text{O})\text{OR}^{15}$ , r)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{NR}^{15}\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ ,  
 s)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{NR}^{15}\text{C}(=\text{NH})\text{NR}^{15}\text{R}^{15}$ , t)  $-\text{O}-\text{C}_{2-6} \text{ alkenyl}-\text{S}(\text{O})_p\text{R}^{15}$ ,  
 u)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{OC}(\text{O})\text{R}^{15}$ , v)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{OC}(\text{O})\text{OR}^{15}$ ,  
 w)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{OC}(\text{O})\text{NR}^{15}\text{R}^{15}$ , x)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ ,  
 y)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{NR}^{15}\text{C}(\text{O})\text{R}^{15}$ , z)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{NR}^{15}\text{C}(\text{O})\text{OR}^{15}$ ,  
 aa)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{NR}^{15}\text{C}(\text{O})\text{NR}^{15}\text{R}^{15}$ ,  
 bb)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{NR}^{15}\text{C}(=\text{NH})\text{NR}^{15}\text{R}^{15}$ , cc)  $-\text{O}-\text{C}_{2-6} \text{ alkynyl}-\text{S}(\text{O})_p\text{R}^{15}$ ; and  
 dd)  $-\text{NR}^{15}\text{R}^{15}$ ;

alternatively, two  $\text{R}^{18}$  groups taken together form  $=\text{O}$ ,  $=\text{NOR}^{15}$ , or  $=\text{NNR}^{15}\text{R}^{15}$ ;  
 $\text{R}^{19}$  is  $\text{R}^{12}$ ;

$\text{R}^{20}$  is selected from the group consisting of:

a)  $\text{R}^{13}$ , b) F, c) Cl, d) Br, e) I, f) CN, g)  $\text{NO}_2$ , and h)  $-\text{OR}^{11}$ ;

alternatively,  $\text{R}^{19}$  and  $\text{R}^{20}$  taken together are  $-\text{O}(\text{CH}_2)_r\text{O}-$ ;

$\text{R}^{21}$ , at each occurrence, independently is selected from the group consisting of:

a) H, b) F, c) Cl, d) Br, e) I, f) CN, g)  $-OR^{11}$ , h)  $NO_2$ , i)  $-NR^{11}R^{11}$ , j)  $C_{1-6}$  alkyl, k)  $C_{1-6}$  acyl, and l)  $C_{1-6}$  alkoxy;

$R^{22}$  is selected from the group consisting of:

- 5 a)  $C_{1-6}$  alkyl, b)  $C_{2-6}$  alkenyl, c)  $C_{2-6}$  alkynyl, d)  $C_{1-6}$  acyl, e)  $C_{1-6}$  alkoxy, f)  $C_{1-6}$  alkylthio, g) saturated, unsaturated, or aromatic  $C_{5-10}$  carbocycle, h) saturated, unsaturated, or aromatic 5-10 membered heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, i)  $-O-C_{1-6}$  alkyl-saturated, unsaturated, or aromatic 5-10 membered heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, j)  $-NR^{11}-C_{1-6}$  alkyl-saturated, 10 unsaturated, or aromatic 5-10 membered heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, k) saturated, unsaturated, or aromatic 10-membered bicyclic ring system optionally containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, l) saturated, unsaturated, or aromatic 13-membered tricyclic ring system optionally containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, m)  $-OR^{11}$ , 15 n)  $-NR^{11}R^{11}$ , o)  $S(O)_rR^{11}$ , and p)  $R^{21}$ ,

20 wherein any of a) - l) optionally is substituted with one or more  $R^{12}$  groups;

alternatively,  $R^{22}$  and one  $R^{21}$  group, taken together with the atoms to which they are bonded, form a 5-7 membered saturated or unsaturated carbocycle, optionally substituted with one or more  $R^{12}$  groups; or a 5-7 membered saturated or unsaturated heterocycle containing one or more atoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally 25 substituted with one or more  $R^{12}$  groups;

$R^{23}$  at each occurrence, independently is selected from the group consisting of:

- a) hydrogen; b) an electron-withdrawing group; c) aryl; d) substituted aryl; e) heteroaryl; f) substituted heteroaryl; and g)  $C_{1-6}$  alkyl, optionally substituted with one or more  $R^{12}$  groups;

30 alternatively, any  $R^{23}$  and any  $R^{20}$ , taken together with the atoms to which they are bonded, form a 5-7 membered saturated or unsaturated carbocycle, optionally substituted with one or more  $R^{12}$  groups; or a 5-7 membered saturated or unsaturated heterocycle containing one



or more atoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally substituted with one or more  $R^{12}$  groups;

p, at each occurrence, is selected from the group consisting of 0, 1, and 2;

r, at each occurrence, is selected from the group consisting of 0, 1, and 2;


5 s, at each occurrence, is selected from the group consisting of 1, 2, 3, or 4;

t, at each occurrence, is selected from the group consisting of 0, 1, or 2;


u, at each occurrence, is selected from the group consisting of 1, 2, 3, 4, or 5; and,





v, at each occurrence, is selected from the group consisting of 0, 1, 2, or 3.


Embodiments of the foregoing compounds include those compounds having the formula:




10











, and











or a pharmaceutically acceptable salt, ester, or prodrug thereof, wherein A, B, n, D, E, R, R<sup>1</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6'</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, and R<sup>10</sup> are as defined hereinabove.

- 5 Other embodiments of the foregoing compounds include those compounds having the formula:

















- 5 or a pharmaceutically acceptable salt, ester, or prodrug thereof, wherein A, B, n, E, R<sup>4</sup>, and R<sup>10</sup> are as defined hereinabove.

Still other embodiments of the foregoing compounds include those compounds having the formula selected from the group consisting of:











- 33 -




, and







or a pharmaceutically acceptable salt, ester, or prodrug thereof, wherein A, B, n, E, and  $R^{10}$  are as defined hereinabove.

- 5 In certain embodiments of the invention, n is 1. In other embodiments,  $A-(B)_n-D$  has the formula  $A-C(O)NH-D$ ,  $A-SO_2NH-D$ , or  $A-C(S)NH-D$ .

Other embodiments of the invention include compounds having the formula:









or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of one or more of the foregoing compounds and a pharmaceutically acceptable carrier. In yet another aspect, the invention provides a method for treating a microbial infection, a fungal infection, a parasitic disease, a proliferative disease, a viral infection, an inflammatory disease, or a gastrointestinal motility disorder in a mammal by administering effective amounts of the compounds of the invention or pharmaceutical compositions of the invention. In embodiments of this aspect, the compounds are administered orally, parentally, or topically. In still another aspect, the invention provides methods for synthesizing any one of the foregoing compounds. In another aspect, the invention provides a medical device, for example, a medical stent, which contains or is coated with one or more of the foregoing compounds.


The invention further provides a family of hybrid antibiotics comprising at least a portion of an heterocyclic side-chain linked via a non-aromatic linker to a macrolide. Exemplary macrolides, linkers, and heterocyclic side-chains useful in the synthesis of the antibiotics include, but are not limited to, the chemical moieties shown below.

#### Macrolides







M4




M5




M6




M7




M8




M9




M10




M11




M12




M13




M14






M15




### Linkers






For the above linker groups, it should be understood that "M" and "O" are included to depict the orientation of the linker group with respect to the other structures that define the compounds of the invention. More specifically, "M" denotes the portion of the compound that includes the macrolide, and "O" denotes the portion of the compound that includes the heterocyclic side-chain.


### Heterocyclic Side-Chains




- 39 -



An exemplary scheme showing the linkage of a macrolide to a heterocyclic side-chain via a linker is set below, where m can be 1, 2, 3, or 4:



5 The above exemplary scheme can be represented as follows:



wherein M is a macrolide selected from the group consisting of M1 through M22, as shown above, B is a linker selected from the group consisting of B1 through B9 as shown above, O is a heterocyclic side chain selected from the group consisting of O1 through O16 as shown above,  
 10 and m is an integer from 1-4.

The various macrolides can be linked via the linkers to the heterocyclic side-chain using conventional chemistries known in the art, such as those discussed above. By using the various combinations of chemical moieties provided, the skilled artisan may synthesize one or more of the exemplary compounds of the present invention. A non-limiting example of a compound that  
 15 one can make based on these exemplary moieties would be wherein the macrolide moiety is selected from M1, the linker moiety is selected from B1 with the variable length chain portion being selected from m = 1, and the heterocyclic side chain moiety is selected from O1. One




skilled in the art would recognize that these illustrated moieties alone can be combined to describe 12,672 unique compounds, i.e. [22 macrolides] x [9 linkers] x [4 chain lengths for each linker] x [16 heterocyclic side chains] = 12, 672.

### 3. Synthesis of the Compounds of the Invention


In another aspect, the invention provides methods for making the compounds of the invention. The following schemes depict some exemplary chemistries available for synthesizing the compounds of the invention. It will be appreciated, however, that the desired compounds may be synthesized using other alternative chemistries known in the art.

The dimethyl amino group of the desosamine sugar of macrolide antibiotics can be monodemethylated to produce the corresponding secondary amine (U.S. Patent No. 3,725,385, Flynn *et al.* (1954) J. AM. CHEM. SOC. 76: 3121; Ku *et al.* (1997) BIOORG. MED. CHEM. LETT. 7: 1203; Stenmark *et al.* (2000) J. ORG. CHEM. 65: 3875). For example, desosamine derivative 1 is available from the degradation of erythromycin. The same chemistry can be employed to produce amine 2 from azithromycin.




Scheme 1 illustrates how amines 1 and 2 can be alkylated with appropriate electrophiles to produce compounds of the present invention that employ linker group L1. Known amine 3 (for a synthesis see: Brickner *et al.* (1996) J. MED. CHEM. 39: 673) was acylated with the appropriate bromoacids to afford bromides 4 and 5. Bromides 4 and 5 were then used to alkylate amine 1 to afford targets 6 and 7, and amine 2 to yield 8 and 9. Compounds of type 6-9 can be converted to thioamide targets of linker type L2 by their reaction with Lawesson's reagent or P<sub>2</sub>S<sub>5</sub>, or many other reagent combinations known to those skilled in the art. It will be understood that a variety of amines similar to 3 could be employed in the chemistry of Scheme 1 (and all subsequent schemes) to provide compounds of the present invention. It is intended that these are within the scope of the present invention.

Scheme 1




- Compounds containing linker group L3 can be made for example by the chemistry shown in Scheme 2. Oxazolidinone amines of type 3 be acylated with commercially available
- 5 bromoacid 10, or the known bromoacid 11 (Bellassoued et al. (1985) TETRAHEDRON 41: 1299), to afford intermediates of type 12 and 13. Alkylation of amines of type 1 and 2 (as non-limiting examples) will provide derivatives of type 14-17.

Scheme 2




Scheme 3 depicts how cis olefin of targets such as **22-25** (with linker group L4) can be made from acetylenic amides of type **20** and **21**. Amines of type **3** can be acylated with the known acetylenic acids **18** (Tsou et al. (2001) J. MED. CHEM. 44: 2719) and **19** (Carey et al. (2001) J. ORG. CHEM. 66: 2526) to afford alkynes **20** and **21** respectively. Hydrogenation of **20** and **21** with hydrogen and Lindlar's catalyst (or other conditions such as diimide reduction) will provide cis olefin intermediates that can be desilylated with tetrabutylammonium fluoride (or acidic conditions such as a mixture of acetic acid in tetrahydrofuran and water) to afford the corresponding alcohols. These alcohols can be converted to tosylates (or other electrophilic species such as a halides) using chemistry well known in the art. These intermediates can serve as alkylating agents for amines such as (but not limited to) **1** and **2** to afford targets **22-25**.

Scheme 3




An alternate approach to targets such as **22-25** is shown in Scheme 4. The siloxy groups of alkynes **20** and **21** can be converted to tosylates to serve as alkylating agents for amines **1** and **2**. The products of the alkylation reaction can then be reduced to give the cis olefin as above to provide targets **22-25**.

Scheme 4



Scheme 5 illustrates how amines such as (but not limited to) 3 can be converted to compounds of the present invention containing linker groups of type L5. Amines 1 and 2 can be alkylated with  $\omega$ -bromo-1-alkanols to afford alcohols such as 26-29. Isocyanates similar to 30 can be made from amines such as 3 using phosgene, carbonyldiimidazole or other reagents well known to those skilled in the art. The reaction of the primary alcohols of compounds 26-29 with isocyanates such as 30 will lead to carbamate targets such as 31-34. It will be understood that amines of type 3 could be converted to isothiocyanates corresponding to 30, and that these intermediates could be used to make thiocarbamate derivatives related to 31-34, containing linker groups of type L6. It will be further appreciated that the primary alcohols of intermediates such as 26-29 could be converted by chemistry well known in the art to primary amines which could then be used to react with isocyanate 30 or the corresponding isothiocyanate to provide urea- and thiourea-linked compounds of the present invention. It is intended that such urea and thiourea derivatives are included in the scope of the present invention.

Scheme 5



#### 4. Characterization of Compounds of the Invention

Compounds designed, selected and/or optimized by methods described above, once produced, may be characterized using a variety of assays known to those skilled in the art to determine whether the compounds have biological activity. For example, the molecules may be characterized by conventional assays, including but not limited to those assays described below, to determine whether they have a predicted activity, binding activity and/or binding specificity.

Furthermore, high-throughput screening may be used to speed up analysis using such assays. As a result, it may be possible to rapidly screen the molecules described herein for activity, for example, as anti-cancer, anti-bacterial, anti-fungal, anti-parasitic or anti-viral agents. Also, it may be possible to assay how the compounds interact with a ribosome or ribosomal subunit and/or are effective as modulators (for example, inhibitors) of protein synthesis using techniques known in the art. General methodologies for performing high-throughput screening are described, for example, in Devlin (1998) High Throughput Screening, Marcel Dekker; and U.S. Patent No. 5,763,263. High-throughput assays can use one or more different assay techniques including, but not limited to, those described below.

(1) *Surface Binding Studies*. A variety of binding assays may be useful in screening new molecules for their binding activity. One approach includes surface plasmon resonance (SPR)

that can be used to evaluate the binding properties of molecules of interest with respect to a ribosome, ribosomal subunit or a fragment thereof.

SPR methodologies measure the interaction between two or more macromolecules in real-time through the generation of a quantum-mechanical surface plasmon. One device, (BIAcore Biosensor RTM from Pharmacia Biosensor, Piscatawy, N.J.) provides a focused beam of polychromatic light to the interface between a gold film (provided as a disposable biosensor "chip") and a buffer compartment that can be regulated by the user. A 100 nm thick "hydrogel" composed of carboxylated dextran that provides a matrix for the covalent immobilization of analytes of interest is attached to the gold film. When the focused light interacts with the free electron cloud of the gold film, plasmon resonance is enhanced. The resulting reflected light is spectrally depleted in wavelengths that optimally evolved the resonance. By separating the reflected polychromatic light into its component wavelengths (by means of a prism), and determining the frequencies that are depleted, the BIAcore establishes an optical interface which accurately reports the behavior of the generated surface plasmon resonance. When designed as above, the plasmon resonance (and thus the depletion spectrum) is sensitive to mass in the evanescent field (which corresponds roughly to the thickness of the hydrogel). If one component of an interacting pair is immobilized to the hydrogel, and the interacting partner is provided through the buffer compartment, the interaction between the two components can be measured in real time based on the accumulation of mass in the evanescent field and its corresponding effects of the plasmon resonance as measured by the depletion spectrum. This system permits rapid and sensitive real-time measurement of the molecular interactions without the need to label either component.

(2) *Fluorescence Polarization.* Fluorescence polarization (FP) is a measurement technique that can readily be applied to protein-protein, protein-ligand, or RNA-ligand interactions in order to derive  $IC_{50}$ s and  $K_d$ s of the association reaction between two molecules. In this technique one of the molecules of interest is conjugated with a fluorophore. This is generally the smaller molecule in the system (in this case, the compound of interest). The sample mixture, containing both the ligand-probe conjugate and the ribosome, ribosomal subunit or fragment thereof, is excited with vertically polarized light. Light is absorbed by the probe fluorophores, and re-emitted a short time later. The degree of polarization of the emitted light is measured. Polarization of the emitted light is dependent on several factors, but most importantly on viscosity of the solution and on the apparent molecular weight of the fluorophore. With proper controls, changes in the degree of polarization of the emitted light depends only on

changes in the apparent molecular weight of the fluorophore, which in-turn depends on whether the probe-ligand conjugate is free in solution, or is bound to a receptor. Binding assays based on FP have a number of important advantages, including the measurement of  $IC_{50}$ s and  $K_d$ s under true homogenous equilibrium conditions, speed of analysis and amenity to automation, and  
5 ability to screen in cloudy suspensions and colored solutions.

(3) *Protein Synthesis*. It is contemplated that, in addition to characterization by the foregoing biochemical assays, the compound of interest may also be characterized as a modulator (for example, an inhibitor of protein synthesis) of the functional activity of the ribosome or ribosomal subunit.

10 Furthermore, more specific protein synthesis inhibition assays may be performed by administering the compound to a whole organism, tissue, organ, organelle, cell, a cellular or subcellular extract, or a purified ribosome preparation and observing its pharmacological and inhibitory properties by determining, for example, its inhibition constant ( $IC_{50}$ ) for inhibiting protein synthesis. Incorporation of  $^3H$  leucine or  $^{35}S$  methionine, or similar experiments can be  
15 performed to investigate protein synthesis activity. A change in the amount or the rate of protein synthesis in the cell in the presence of a molecule of interest indicates that the molecule is a modulator of protein synthesis. A decrease in the rate or the amount of protein synthesis indicates that the molecule is a inhibitor of protein synthesis.

Furthermore, the compounds may be assayed for anti-proliferative or anti-infective  
20 properties on a cellular level. For example, where the target organism is a microorganism, the activity of compounds of interest may be assayed by growing the microorganisms of interest in media either containing or lacking the compound. Growth inhibition may be indicative that the molecule may be acting as a protein synthesis inhibitor. More specifically, the activity of the compounds of interest against bacterial pathogens may be demonstrated by the ability of the  
25 compound to inhibit growth of defined strains of human pathogens. For this purpose, a panel of bacterial strains can be assembled to include a variety of target pathogenic species, some containing resistance mechanisms that have been characterized. Use of such a panel of organisms permits the determination of structure-activity relationships not only in regards to potency and spectrum, but also with a view to obviating resistance mechanisms. The assays may  
30 be performed in microtiter trays according to conventional methodologies as published by The National Committee for Clinical Laboratory Standards (NCCLS) guidelines (NCCLS. M7-A5- Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard-Fifth Edition. NCCLS Document M100-S12/M7 (ISBN 1-56238-394-9)).



## 5. Formulation and Administration

The compounds of the invention may be useful in the prevention or treatment of a variety of human or other animal disorders, including for example, bacterial infection, fungal infections, viral infections, parasitic diseases, and cancer. It is contemplated that, once identified, the active molecules of the invention may be incorporated into any suitable carrier prior to use. The dose of active molecule, mode of administration and use of suitable carrier will depend upon the intended recipient and target organism. The formulations, both for veterinary and for human medical use, of compounds according to the present invention typically include such compounds in association with a pharmaceutically acceptable carrier.

The carrier(s) should be "acceptable" in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient. Pharmaceutically acceptable carriers, in this regard, are intended to include any and all solvents, dispersion media, coatings, anti-bacterial and anti-fungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds (identified or designed according to the invention and/or known in the art) also can be incorporated into the compositions. The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy/microbiology. In general, some formulations are prepared by bringing the compound into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation.

A pharmaceutical composition of the invention should be formulated to be compatible with its intended route of administration. Examples of routes of administration include oral or parenteral, for example, intravenous, intradermal, inhalation, transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the

adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide.

Useful solutions for oral or parenteral administration can be prepared by any of the methods well known in the pharmaceutical art, described, for example, in Remington's  
5 Pharmaceutical Sciences, (Gennaro, A., ed.), Mack Pub., (1990). Formulations for parenteral administration can also include glycocholate for buccal administration, methoxysalicylate for rectal administration, or citric acid for vaginal administration. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Suppositories for rectal administration also can be prepared by mixing the drug with a non-  
10 irritating excipient such as cocoa butter, other glycerides, or other compositions which are solid at room temperature and liquid at body temperatures. Formulations also can include, for example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, and hydrogenated naphthalenes. Formulations for direct administration can include glycerol and other compositions of high viscosity. Other potentially useful parenteral carriers for these drugs  
15 include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration can contain as excipients, for example, lactose, or can be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Retention enemas also can be used for rectal  
20 delivery.

Formulations of the present invention suitable for oral administration may be in the form of: discrete units such as capsules, gelatin capsules, sachets, tablets, troches, or lozenges, each containing a predetermined amount of the drug; a powder or granular composition; a solution or a suspension in an aqueous liquid or non-aqueous liquid; or an oil-in-water emulsion or a water-in-  
25 oil emulsion. The drug may also be administered in the form of a bolus, electuary or paste. A tablet may be made by compressing or moulding the drug optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the drug in a free-flowing form such as a powder or granules, optionally mixed by a binder, lubricant, inert diluent, surface active or dispersing agent. Moulded tablets may be made by  
30 moulding, in a suitable machine, a mixture of the powdered drug and suitable carrier moistened with an inert liquid diluent.

Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients.

Oral compositions prepared using a fluid carrier for use as a mouthwash include the compound in the fluid carrier and are applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the

5 following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose; a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

10       Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor ELTM (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). It should be stable under the conditions of manufacture and  
15 storage and should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of  
20 dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

25       Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile  
30 powders for the preparation of sterile injectable solutions, methods of preparation include vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Formulations suitable for intra-articular administration may be in the form of a sterile aqueous preparation of the drug that may be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems may also be used to present the drug for both intra-articular and ophthalmic administration.

Formulations suitable for topical administration, including eye treatment, include liquid or semi-liquid preparations such as liniments, lotions, gels, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops. Formulations for topical administration to the skin surface can be prepared by dispersing the drug with a dermatologically acceptable carrier such as a lotion, cream, ointment or soap. Particularly useful are carriers capable of forming a film or layer over the skin to localize application and inhibit removal. For topical administration to internal tissue surfaces, the agent can be dispersed in a liquid tissue adhesive or other substance known to enhance adsorption to a tissue surface. For example, hydroxypropylcellulose or fibrinogen/thrombin solutions can be used to advantage. Alternatively, tissue-coating solutions, such as pectin-containing formulations can be used.

For inhalation treatments, inhalation of powder (self-propelling or spray formulations) dispensed with a spray can, a nebulizer, or an atomizer can be used. Such formulations can be in the form of a fine powder for pulmonary administration from a powder inhalation device or self-propelling powder-dispensing formulations. In the case of self-propelling solution and spray formulations, the effect may be achieved either by choice of a valve having the desired spray characteristics (*i.e.*, being capable of producing a spray having the desired particle size) or by incorporating the active ingredient as a suspended powder in controlled particle size. For administration by inhalation, the compounds also can be delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration also can be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants generally are known in the art, and include, for example, for transmucosal administration, detergents and bile salts. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds typically are formulated into ointments, salves, gels, or creams as generally known in the art.

The active compounds may be prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. Liposomal suspensions can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

Oral or parenteral compositions can be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals. Furthermore, administration can be by periodic injections of a bolus, or can be made more continuous by intravenous, intramuscular or intraperitoneal administration from an external reservoir (*e.g.*, an intravenous bag).

Where adhesion to a tissue surface is desired the composition can include the drug dispersed in a fibrinogen-thrombin composition or other bioadhesive. The compound then can be painted, sprayed or otherwise applied to the desired tissue surface. Alternatively, the drugs can be formulated for parenteral or oral administration to humans or other mammals, for example, in therapeutically effective amounts, *e.g.*, amounts that provide appropriate concentrations of the drug to target tissue for a time sufficient to induce the desired effect.

Where the active compound is to be used as part of a transplant procedure, it can be provided to the living tissue or organ to be transplanted prior to removal of tissue or organ from the donor. The compound can be provided to the donor host. Alternatively or, in addition, once removed from the donor, the organ or living tissue can be placed in a preservation solution containing the active compound. In all cases, the active compound can be administered directly to the desired tissue, as by injection to the tissue, or it can be provided systemically, either by oral or parenteral administration, using any of the methods and formulations described herein and/or known in the art. Where the drug comprises part of a tissue or organ preservation

solution, any commercially available preservation solution can be used to advantage. For example, useful solutions known in the art include Collins solution, Wisconsin solution, Belzer solution, Eurocollins solution and lactated Ringer's solution.

Active compound as identified or designed by the methods described herein can be administered to individuals to treat disorders (prophylactically or therapeutically). In conjunction with such treatment, pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a drug as well as tailoring the dosage and/or therapeutic regimen of treatment with the drug.

In therapeutic use for treating, or combating, bacterial infections in mammals, the compounds or pharmaceutical compositions thereof will be administered orally, parenterally and/or topically at a dosage to obtain and maintain a concentration, that is, an amount, or blood-level or tissue level of active component in the animal undergoing treatment which will be antimicrobially effective. The term "effective amount" is understood to mean that the compound of the invention is present in or on the recipient in an amount sufficient to elicit biological activity, for example, anti-microbial activity, anti-fungal activity, anti-viral activity, anti-parasitic activity, and/or anti-proliferative activity. Generally, an effective amount of dosage of active component will be in the range of from about 0.1 to about 100, more preferably from about 1.0 to about 50 mg/kg of body weight/day. The amount administered will also likely depend on such variables as the type and extent of disease or indication to be treated, the overall health status of the particular patient, the relative biological efficacy of the compound delivered, the formulation of the drug, the presence and types of excipients in the formulation, and the route of administration. Also, it is to be understood that the initial dosage administered may be increased beyond the above upper level in order to rapidly achieve the desired blood-level or tissue level, or the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation. If desired, the daily dose may also be divided into multiple doses for administration, for example, two to four times per day.



## 6. Examples





In the following examples, nuclear magnetic resonance (NMR) spectra were obtained on a Bruker Avance 300 or Avance 500 spectrometer, or in some cases a GE-Nicolet 300 spectrometer. Common reaction solvents were either high performance liquid chromatography (HPLC) grade or American Chemical Society (ACS) grade, and anhydrous as obtained from the manufacturer unless otherwise noted. "Chromatography" or "purified by silica gel" refers to flash column chromatography using silica gel (EM Merck, Silica Gel 60, 230-400 mesh) unless otherwise noted.

### Example 1 - Exemplary Compounds





Exemplary compounds synthesized in accordance with the invention are listed in Table 1.

TABLE 1

| Compound Number | Structure  |
|-----------------|--|
| 6               |  |
| 7               |  |

|    |  |
|----|--|
| 8  |    |
| 9  |    |
| 35 |   |
| 37 |  |



|     |  |
|-----|--|
| 38  |    |
| 39  |    |
| 102 |   |
| 103 |  |

**Example 2 - Synthesis of Compounds 6-9****Synthesis of amine 2**

Azithromycin (0.80 g, 1.02 mmol) and sodium acetate (NaOAc) (0.712 g, 8.06 mmol) were dissolved in 80% aqueous methanol (MeOH) (25 mL). The solution was kept at 50°C followed by addition of iodine (I<sub>2</sub>) (0.272 g, 1.07 mmol) in three batches within 3 minutes (min). The reaction was maintained at a pH between 8 – 9 by adding 1N sodium hydroxide (NaOH) (1 mL) at 10 min and 45 min intervals. The solution turned colorless within 45 min, however, stirring was continued for 2 hours (hr). TLC (methylene chloride/methanol/ammonium hydroxide (CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH) 10:1:0.05) after 2 hr showed a single major product (R<sub>f</sub> = 0.66). The reaction was cooled to room temperature (RT), poured into water (H<sub>2</sub>O) (75 mL) containing NH<sub>4</sub>OH (1.5 mL) and extracted with chloroform (CHCl<sub>3</sub>) (3 x 30 mL). The combined organic layer was washed with H<sub>2</sub>O (30 mL) containing NH<sub>4</sub>OH (1.5 mL), dried over sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated to give a white residue. The crude was purified on silica gel column eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH 18:1:0.05 to 10:1:0.05 to provide amine 2 (0.41 g, 55% yield).

**Synthesis of bromide 4**

To a solution of amine 3 (0.43g, 1.46 mmol) in anhydrous tetrahydrofuran (THF) (10 mL) was added 3-bromopropionic acid (0.23g, 1.5 mmol) and 1,3-dicyclohexylcarbodiimide (DCC) (0.36g, 1.72 mmol). A precipitate developed within 2 min. Stirring was continued at RT for 2 hr, and the reaction was then filtered to remove the precipitated urea. CH<sub>2</sub>Cl<sub>2</sub> (60 mL) and saturated ammonium chloride (NH<sub>4</sub>Cl) were added to the filtrate. The two layers were separated, and the organic layer was washed with saturated NH<sub>4</sub>Cl (2 x 30 mL) then saturated brine (1 x 30 mL), and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated, and the residue was purified on a silica gel column eluting with ethyl acetate (EtOAc)/Hexanes 8:1 to 9:1 to give 4 as a white solid (0.42 g, 67% yield). <sup>1</sup>H-NMR (300 MHz, deuterated chloroform (CDCl<sub>3</sub>); partial): δ 7.30 (dd, *J* = 14, 3 Hz, 1H), 6.95 (dd, *J* = 9, 2 Hz, 1H), 6.78 (t, *J* = 9 Hz, 1H), 6.19 (t, *J* = 6 Hz, 1H), 4.67 (m, 1H), 3.45-3.89 (m, 10H), 2.92 (t, *J* = 5 Hz, 4H), 2.65 (m, 2H).

**Synthesis of bromide 5**

Compound 5 was synthesized from amine 3 and 4-bromobutyric acid as described for amide 4. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial): δ 7.36 (dd, *J* = 14, 3 Hz, 1H), 7.01 (dd, *J* = 9, 2

Hz, 1H), 6.85 (t,  $J = 9$  Hz, 1H), 5.95 (t,  $J = 6$  Hz, 1H), 4.69 (m, 1H), 3.79 (t,  $J = 5$  Hz, 4H), 3.57-3.70 (m, 4H), 3.33 (t,  $J = 6$  Hz, 2H), 2.92 (t,  $J = 5$  Hz, 4H), 2.32 (m, 2H), 2.05 (m, 2H).

### Synthesis of compound 8

A mixture of amine 2 (0.128 g, 0.174 mmol) and bromide 4 (0.12 g, 0.28 mmol) in anhydrous THF (15 mL) and *N,N*-diisopropylethylamine (Hunig's base or *i*-Pr<sub>2</sub>NEt) (3 mL) was heated under reflux overnight. The reaction was poured into CH<sub>2</sub>Cl<sub>2</sub> (60 mL), washed with saturated NH<sub>4</sub>Cl (3 x 30 mL) then saturated brine (1 x 30 mL), and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated, and the crude material was purified on silica gel column first eluting with 2-6% MeOH in CH<sub>2</sub>Cl<sub>2</sub> then with CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH 15:1: 0.05 to 12:1:0.05 to 10:1 to give compound 8 (0.0392 g, 21% yield). <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial):  $\delta$  9.01 (bs, 1H), 7.46 (dd,  $J = 14$ , 2 Hz, 1H), 7.05 (dd,  $J = 9$ , 2 Hz, 1H), 6.91 (t,  $J = 9$  Hz, 1H), 5.10 (bs, 1H), 4.72 (m, 3H), 4.41 (d,  $J = 7$  Hz, 1H), 4.26 (bs, 1H), 3.99 (m, 2H), 3.86 (t,  $J = 5$  Hz, 4H), 3.05 (t,  $J = 5$  Hz, 4H).

### Synthesis of compound 9

Compound 9 was made from amine 2 and bromide 5 as described above for compound 8. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial):  $\delta$  8.99 (bs, 1H), 7.40 (dd,  $J = 14$ , 3 Hz, 1H), 7.05 (dd,  $J = 9$ , 2 Hz, 1H), 6.87 (t,  $J = 9$  Hz, 1H), 5.02 (d,  $J = 2$  Hz, 1H), 4.85 (m, 1H), 4.69 (m, 1H), 4.34 (d,  $J = 7$  Hz, 1H), 3.82 (t,  $J = 4$  Hz, 4H), 3.01 (t,  $J = 4$  Hz, 4H).

### Synthesis of compound 6

Compound 6 was made from amine 1 and amide 4 as described above for compound 8. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial):  $\delta$  8.97 (t,  $J = 6$  Hz, 1H), 7.39 (dd,  $J = 14$ , 3 Hz, 1H), 7.04 (dd,  $J = 9$ , 2 Hz, 1H), 6.87 (t,  $J = 9$  Hz, 1H), 5.01 (d,  $J = 2$  Hz, 1H), 4.84 (d,  $J = 4$  Hz, 1H), 4.69 (m, 1H), 4.33 (d,  $J = 7$  Hz, 1H), 3.82 (t,  $J = 4$  Hz, 4H), 3.01 (t,  $J = 5$  Hz, 4H).

### Synthesis of compound 7

Compound 7 was made from amine 1 and amide 5 as described above for compound 8. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial):  $\delta$  8.99 (bs, 1H), 7.40 (dd,  $J = 14$ , 3 Hz, 1H), 7.05 (dd,  $J = 9$ , 2 Hz, 1H), 6.87 (t,  $J = 9$  Hz, 1H), 5.02 (d,  $J = 3$  Hz, 1H), 4.85 (m, 1H), 4.69 (m, 1H), 4.34 (d,  $J = 7$  Hz, 1H), 3.82 (t,  $J = 4$  Hz, 4H), 3.01 (t,  $J = 4$  Hz, 4H).

### Example 3 - Synthesis of Compound 35

### Synthesis of bromide 36

Compound **36** was synthesized from amine **3** and bromoacetic acid as described for amide **4**. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial): δ 7.36 (dd, *J* = 11, 2 Hz, 1H), 7.02 (m, 2H), 6.93 (dd, *J* = 9, 2 Hz, 1H), 4.68 (m, 1H), 3.93-3.53 (m, 10H), 3.01 (t, *J* = 5 Hz, 4H). LCMS (ESI) *m/z* 418 (M+H<sup>+</sup>).


### Synthesis of compound 35

Compound **35** was made from amine **2** and amide **36** as described for compound **8**. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial): δ 8.25, (t, *J* = 6 Hz, 1H), 7.40 (dd, *J* = 14, 3 Hz, 1H), 6.99 (dd, *J* = 9, 2 Hz, 1H), 6.84 (t, *J* = 9 Hz, 1H), 4.96 (d, *J* = 4 Hz, 1H), 4.69 (m, 2H), 4.31 (d, *J* = 8 Hz, 2H), 4.19 (bs, 1H), 3.80 (t, *J* = 5 Hz, 4H), 2.98 (t, *J* = 5 Hz, 4H). LCMS (ESI) *m/z* 1070 (M+H<sup>+</sup>).

### Example 4 - Synthesis of Sulfonamide 37

Scheme 6 shows the synthesis of sulfonamide 37. The hydrochloride salt (41) of the known amine (see: Barbachyn, M. *et al.* International Patent Application WO95/07271) was converted to sulfonamide 42. Sulfonamide 42 was used to alkylate amine 171 to afford sulfonamide 37.

Scheme 6



## 20 Synthesis of sulfonamide 42

To a stirred solution of amine salt **41** (332 mg, 1.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added sequentially triethylamine (Et<sub>3</sub>N) (0.5 mL), and 2-chloro-ethanesulfonyl chloride

(ClCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Cl) (163 mg, 1.0 mmol). The reaction mixture was stirred at (RT) for 16 hr, then diluted to 50 mL with CH<sub>2</sub>Cl<sub>2</sub> and washed with saturated aqueous sodium bicarbonate (NaHCO<sub>3</sub>) (50 mL) and brine (25 mL). The organic fraction was dried over potassium carbonate (K<sub>2</sub>CO<sub>3</sub>), filtered and concentrated to give 490 mg of a yellow solid which was purified by silica gel chromatography (15 mm x 6" column eluted with 1:1 acetone/ hexanes) to afford sulfonamide **42** as a colorless solid (380 mg, 0.90 mmol) which was used as-is for the next reaction.


### Synthesis of sulfonamide 37

To a stirred solution of amine **2** (0.10 g, 0.14 mmol) in 3 mL **i**-Pr<sub>2</sub>NEt was added **42** (0.10 g, 0.24 mmol). The solution was sealed in a 1 dram vial and heated in a 100 °C oil bath for 16 h. The reaction mixture was cooled, then diluted to 50 mL with CH<sub>2</sub>Cl<sub>2</sub> and washed with saturated aqueous NaHCO<sub>3</sub> (50 mL) and brine (25 mL). The organic fraction was dried over K<sub>2</sub>CO<sub>3</sub>, filtered and concentrated to give 170 mg of a yellow oil which was purified by silica gel chromatography (15 mm x 6" column eluted with 33:1 CH<sub>2</sub>Cl<sub>2</sub> / 2N NH<sub>3</sub> in MeOH) to afford sulfonamide **37** as a white solid (105 mg, 0.094 mmol). <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>; partial): δ 7.46 (dd, *J* = 14, 2 Hz, 1H), 7.06 (t, *J* = 6 Hz, 1H), 6.97 (dd, *J* = 9, 2 Hz, 1H), 6.84 (t, *J* = 9 Hz, 1H), 4.83 (d, *J* = 2 Hz, 1H), 4.58 (d, *J* = 8 Hz, 1H), 4.38 (d, *J* = 7 Hz, 1H), 4.15 (d, *J* = 6 Hz, 1H), (3.22, s, 3H), 2.23 (s, 3H), 2.31 (s, 3H), 1.34-1.27 (m, 8H), 1.27-1.15 (m, 10H), 1.10-0.99 (m, 9H), 0.89-0.78 (m, 6H). LCMS (ESI) *m/z* 1120 (M + H)<sup>+</sup>, 560.8 (M + 2H)<sup>2+</sup>.

### Example 5 - Synthesis of Compound 38

20 Scheme 7 depicts the synthesis of thioamide **38**. Bromide **4** was converted to a mixture of thioamides **43** and **44**, which was subsequently used to alkylate amine **2** to afford thioamide **38**.

Scheme 7



**Synthesis of thioamides 43 and 44**

Bromopropionamide 4 (0.190 g, 0.440 mmol) and Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide) (0.092 g, 0.220 mmol) were dissolved in anhydrous THF (10 mL) and the solution was heated under reflux for 2 h. The solvent was evaporated; and the crude was purified on a silica gel column, eluting with EtOAc/Hexanes (1:1 to 2:1) to give a brown-white solid. LCMS analysis showed that the isolated product was approximately a 7:3 mixture of thioamide 43 [ $m/e$  448.1 ( $M+H$ )<sup>+</sup>] and thioamide 44 [ $m/e$  366.0 ( $M+H$ )<sup>+</sup>] respectively (0.072 g, combined yield).


**Synthesis of thioamide 38**

A mixture of amine 2 (0.250 g, 0.323 mmol) and thioamides 43 and 44 (0.072 g, 0.160 mmol based on 43) in anhydrous THF (5 mL) and Hunig's base (5 mL) was heated at 100°C for 12 h. The reaction was poured into CH<sub>2</sub>Cl<sub>2</sub> (60 mL), extracted with a mixture containing saturated NH<sub>4</sub>Cl/NH<sub>4</sub>OH (pH = 9.5, 1 x 30 mL) and saturated brine (1 x 30 mL) and the organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated and the crude was purified on a silica gel column first eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (18:1) then with CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH (18:1: 0.03 to 18:1:0.05 to 15:1:0.05) to give 38 as a white solid (0.027 g, 15% yield). <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial):  $\delta$  11.56 (bs, 1H), 7.40 (dd,  $J$  = 14, 2 Hz, 1H), 6.95 (dd,  $J$  = 9, 3 Hz, 1H), 6.83 (t,  $J$  = 9 Hz, 1H), 4.93 (m, 2H), 4.61 (m, 1H), 4.33 (d,  $J$  = 7 Hz, 1H), 3.78 (t,  $J$  = 5 Hz, 4H), 2.96 (t,  $J$  = 5 Hz, 4H). LCMS (ESI)  $m/z$  1100.4 ( $M+H$ )<sup>+</sup>.

**Example 6 - Synthesis of Compound 39**

Scheme 8 illustrates the synthesis of amine 39. Amine 45 was alkylated with 3-bromopropanol, and the resulting diol 46 was converted to monotosylate 47. Alkylation of amine 2 with tosylate 47 yielded amine 39.

Scheme 8



**Synthesis of diol 46**

A mixture of amine 45 (0.50g, 2.38 mmol; synthesized using the chemistry reported in the literature (azide synthesized via Brickner, S.J. *et al. J. Med. Chem.* 1996, 39, 673; then reduction of the azide to the amine) and 3-bromopropanol (0.42 mL, 4.68 mmol) in THF (10 mL) and *i*-Pr<sub>2</sub>NEt (10 mL) was heated at 100°C for 48 h. The solvent was evaporated and the crude was purified on a silica gel column, eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (14:1 to 12:1) to give alcohol 46 as thick oil (0.71 g, 91% yield). LCMS (ESI) *m/z* 326.9 (M+H)<sup>+</sup>.

**Synthesis of tosylate 47**

To a solution of alcohol 46 (0.35 g, 1.07 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and Et<sub>3</sub>N (0.19 mL, 1.30 mmol) was added *p*-toluenesulfonyl chloride (TsCl) (0.21 g, 1.07 mmol) at 0°C. The reaction was allowed to warm to RT and stirred for 20 hr. The solvent was evaporated and the crude was purified on a silica gel column, eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (20:1 to 18:1 to 16:1) to give tosylate 47 (0.096 g, 19% yield). LCMS (ESI) *m/z* 481.0 (M+H)<sup>+</sup>.

**Synthesis of amine 39**


A mixture of amine 2 (0.250 g, 0.323 mmol) and tosylate 47 (0.092 g, 0.192 mmol) in anhydrous THF (5 mL) and *i*-Pr<sub>2</sub>NEt (5 mL) was heated at 100°C for 24 hr. The solvent was evaporated and the crude was purified on a silica gel column eluting with CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH (18:1: 0.04 to 16:1:0.04 to 14:1:0.05 to 12:1:0.05) to give amine 39 as a white solid (0.050 g, 25% yield). <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>; partial): δ 7.37 (m, 1H), 7.25 (m, 1H), 7.18 (m, 1H), 6.77 (m, 1H), 4.94 (d, *J* = 4 Hz, 1H), 4.73 (m, 1H), 4.60 (d, *J* = 9 Hz, 1H), 4.37 (d, *J* = 7 Hz, 1H), 4.20 (d, *J* = 4 Hz, 4H), 3.69 (m, 3H), 0.83 (m, 6H). LCMS (ESI) *m/z* 1043.6 (M+H)<sup>+</sup>.

**Example 7 - Synthesis of Compound 102**

Scheme 9 illustrates the synthesis of compound 102. Amine 45 was condensed with succinic acid monomethyl ester to provide amide ester 110, which was hydrolyzed to carboxylic acid 111. This carboxylic acid 111 was condensed with 4'-α-hydroxy-azithromycin 114, synthesis provided in Example 9, to yield compound 102.

- 63 -

Scheme 9

**Synthesis of amide ester 110**

To a solution of amine **45** (50mg, 0.24 mmol), and succinic acid monomethyl ester (47mg, 0.36 mmol) in 2 mL CH<sub>2</sub>Cl<sub>2</sub>, was added 74 mg DCC (0.36 mmol) and the mixture stirred at RT for 4 hr. The reaction mixture was placed directly on a silica gel column and eluted with hexane/acetone (3:1) to afford amide ester **110** (80 mg) as a white solid.

**Synthesis of carboxylic acid 111**

Amide ester **110** was dissolved in 5 mL ethanol (EtOH) and 0.5 mL 2N potassium hydroxide (KOH) (aq) was added. After stirring for 4 hr at RT the reaction mixture was diluted to 50 mL with 0.1N hydrochloric acid (HCl) (aq) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 30 mL). The combined organic fractions were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford carboxylic acid **111** as a white solid.

**Synthesis of compound 102**


To a stirred solution of this carboxylic acid **111** (24 mg, 0.07 mmol) and 4'- $\alpha$ -hydroxy azithromycin **2** (38 mg, 0.05 mmol) in 0.5 mL CH<sub>2</sub>Cl<sub>2</sub> was added 76  $\mu$ L of a 20% w/v solution of DCC in CH<sub>2</sub>Cl<sub>2</sub>. The mixture was stirred at RT for 15 hr then the entire reaction mixture was placed directly on a silica gel flash column and eluted with 1:20:0.1 MeOH/CH<sub>2</sub>Cl<sub>2</sub>/NH<sub>3</sub> to afford compound **102** as a white solid (41 mg, 0.04 mmol). LCMS (ESI)  $m/z$  529.5 (M + 2H)<sup>2+</sup>.



**Example 8 - Synthesis of Compound 103**

Scheme 10 illustrates the synthesis of compound 103. Amine 45 was condensed with Boc-L-glutamic acid 5-benzyl ester to provide amide ester 112, which was hydrolyzed to carboxylic acid 113. This carboxylic acid 113 was condensed with amine 114 to yield compound 103.

Scheme 10

**Synthesis of amide ester 112**

To a solution of amine 45 (50mg, 0.24 mmol), and Boc-L- glutamic acid-5-benzyl ester (97mg, 0.29 mmol) in 2 mL CH<sub>2</sub>Cl<sub>2</sub>, was added 74 mg DCC (0.36 mmol) and the mixture stirred at RT for 4 hr. The reaction mixture was placed directly on a silica gel column and eluted with hexane/acetone (4:1) to afford amide ester 112 (138 mg) as a white solid.

**Synthesis of carboxylic acid 113**

Amide ester 112 was dissolved in 5 mL EtOH and 0.5 mL 2N KOH(aq) was added. After stirring for 4 hr at RT the reaction mixture was diluted to 50 mL with 0.1N HCl(aq) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 30 mL). The combined organic fractions were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford carboxylic acid 113 as a white solid (129 mg).


**Synthesis of compound 103**

To a stirred solution of this carboxylic acid **113** (33 mg, 0.08 mmol) and 4'- $\alpha$ -hydroxy azithromycin **2** (38 mg, 0.05 mmol) in 0.5 mL CH<sub>2</sub>Cl<sub>2</sub> was added 76  $\mu$ L of a 20% w/v solution of DCC in CH<sub>2</sub>Cl<sub>2</sub>. The mixture was stirred at RT for 15 hr then the entire reaction mixture was placed directly on a silica gel flash column and eluted with 1:20:0.1 MeOH/CH<sub>2</sub>Cl<sub>2</sub>/NH<sub>3</sub> to afford compound **103** as a white solid (41 mg, 0.04 mmol). LCMS (ESI)  $m/z$  544.0 (M-Boc + 2H)<sup>2+</sup>.

**Example 9 - Synthesis of Compound 114**

Scheme 11 illustrates the synthesis of 4'- $\alpha$ -hydroxy-azithromycin **114** from azithromycin **115** via amine oxide **116**. The amine oxide **116** is converted to alkenyl compound **117**, which is converted to epoxide N-oxide **118**, which is subsequently converted to azido alcohol **119**, and then to compound **114**.

Scheme 11

**15 Synthesis of azithromycin-3'-N-oxide 116**

Azithromycin **115** (50 g, 66.8 mmol) was dissolved in enough warm acetone to make 150 mL of solution. This solution was allowed to cool to ambient temperature prior to addition of 40 ml of 30% w/w aqueous H<sub>2</sub>O<sub>2</sub>. Following a mild exotherm, the solution was allowed to cool to ambient temperature and stirred for 3.5 hr. The reaction mixture was diluted to 2 L with

CH<sub>2</sub>Cl<sub>2</sub> and the resulting gelatinous mixture was stirred vigorously for 1 hr to afford a cloudy suspension. This suspension was washed with a 5:1 mixture of saturated aqueous NaHCO<sub>3</sub> and 10% w/v aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (2 x 600 mL) and with brine (1 x 800 mL). The aqueous washes were combined and adjusted to pH 12 with 2N KOH and then further extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 300 mL). The combined organic extracts were dried over K<sub>2</sub>CO<sub>3</sub>, filtered, and concentrated *in vacuo*. As the volume of the extracts was reduced crystals began to form; when the total volume of the extracts had been reduced to 700 mL the solution was placed in a stoppered flask and stored at RT overnight. The solids were collected by vacuum filtration, rinsed with cold ether, and dried under vacuum to afford 34 g of white needle-like crystals. The filtrate was treated as before to yield two additional crops of crystalline product **201** for a total yield of 51 g (66.7 mmol 99%). LCMS (ESI) *m/z* 765.6 (M + H)<sup>+</sup>.

#### Synthesis of 3'-desdimethylamino-4'-dehydro-azithromycin (alkenyl compound) **117**

A 300 mL pear-shaped recovery flask was charged with Azithromycin-3'-N-oxide **116** (35 g, 45.8 mmol) and placed on a rotary evaporator. The pressure was reduced to 0.5 torr and the flask was rotated slowly in an oil bath while the temperature was gradually increased to 175 °C. The mixture was held under vacuum at this temperature for 1.5 hr then cooled to RT and flushed with argon. The resulting tan solid was dissolved in 800 mL of boiling acetonitrile (CH<sub>3</sub>CN). The solution was allowed to cool slowly to RT and then placed in a -20 °C freezer overnight. The solids were collected by vacuum filtration and washed with cold acetonitrile to afford 19.1 g of **117** as off-white crystals. The filtrate was concentrated and the residue treated as above to afford two additional crops of **117** product for a total yield of 27.7 g (39.4 mmol, 86%). LCMS (ESI) *m/z* 704.5 (M + H)<sup>+</sup>.

#### Synthesis of 3'-desdimethylamino-4'-dehydro-3',4'-epoxy-9'-N-oxo-azithromycin (epoxide N-oxide) **118**

To a methanol solution of **117** (25.0g, 35.5 mmol in 100 mL) was added *meta*-chloroperbenzoic acid (20.4g, 89 mmol). The reaction mixture was stirred at RT for 14 hr at which time an additional 10 g portion of *m*CPBA was added. The solution was stirred for an additional 4 hr, then diluted with 1200 mL CH<sub>2</sub>Cl<sub>2</sub> and washed with saturated aqueous NaHCO<sub>3</sub> (2 x 500 mL) and brine (1 x 500 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 500 mL). The combined organic extracts were dried on K<sub>2</sub>CO<sub>3</sub>, filtered, and concentrated to give a white foam (30.7g) which was purified by silica gel chromatography (125mm x 6")

column eluted with 7.5% 2N ammonia (NH<sub>3</sub>) in MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) to afford compound **118** as a white solid (25.7 g, 35.0 mmol, 98%). LCMS (ESI) *m/z* 779.6 (M + H)<sup>+</sup>.

**Synthesis of 3'β-azido-4'α-hydroxy-9'N-oxo-3'-desdimethylamino-azithromycin (azido alcohol) 119**

5 Epoxide N-oxide **118** (20.0g, 27.2 mmol) was dissolved in 88 mL of 10:1 dimethylsulfoxide-water (DMSO-H<sub>2</sub>O) to which was added sodium azide (NaN<sub>3</sub>) (17.7g, 270 mmol) and magnesium perchlorate octahydrate (Mg(ClO<sub>4</sub>)•8H<sub>2</sub>O) (13.5g, 40.8 mmol). The mixture was stirred under argon at 85 °C for 16 hr then cooled to RT and poured into saturated aqueous NaHCO<sub>3</sub> (1L) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (5 x 500 mL). The combined organic extracts  
10 were dried over K<sub>2</sub>CO<sub>3</sub>, filtered, and concentrated to afford a white foam (29 g). This material was dissolved in hot CH<sub>3</sub>CN (1.2L) and allowed to sit overnight at RT. The solids were filtered from the solution and rinsed with additional CH<sub>3</sub>CN. The 8.7 g of crystalline solid thus obtained was confirmed by NMR and x-ray analysis to be pure 3'α-hydroxy-4'β-azido-9'N-oxo-3'-desdimethylamino-azithromycin formed by addition of the azide at the 4' carbon of the epoxide.  
15 The mother liquors were concentrated and the residue again dissolved in boiling CH<sub>3</sub>CN from which a second 3.0 g crop of the undesired isomer was obtained in pure form. The mother liquors, now enriched in the product **119**, were concentrated and the residue purified by silica gel chromatography (50 mm x 8" column eluted with 0-8% 2N NH<sub>3</sub> in MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) to afford an additional 2.9 g of the earlier-eluting 4'β-azide along with compound **119** (6.5 g, 8.3 mmol, 31  
20 %). LCMS (ESI) *m/z* 736.6 (M + H)<sup>+</sup>.

**Synthesis of 4'α-hydroxy-azithromycin 114**

A heavy-walled pressure tube was charged with an ethanol solution of **119** (1.73 g, 2.22 mmol in 20 mL) and 20% palladium (Pd) on charcoal (C) (0.14 g containing 50% H<sub>2</sub>O). The reaction mixture was stirred under an hydrogen (H<sub>2</sub>) atmosphere (15 psig) at RT for 14 hr at  
25 which time 2 mL 37% aqueous formaldehyde (CH<sub>2</sub>O), 1 mL formic acid (HCO<sub>2</sub>H), and an additional 50 mg Pd on C were added. The hydrogen pressure was increased to 30 psig and stirring was continued for 24 hr, at which time an additional 100 mg charge of Pd was added and the H<sub>2</sub> pressure was increased to 90 psig. After an additional 24 hr at this pressure the reaction mixture was purged with argon, filtered, diluted with 100 mL toluene, and concentrated *in vacuo*  
30 to afford 1.9 g of a colorless glass. The crude product was purified by silica gel chromatography

(25 mm x 6" column eluted with 7% 2N NH<sub>3</sub> in MeOH/ CH<sub>2</sub>Cl<sub>2</sub>) to afford compound 114 as a white solid (0.78 g, 1.0 mmol, 45%). LCMS (ESI) *m/z* 765.5 (M + H)<sup>+</sup>.

#### INCORPORATION BY REFERENCE

The entire disclosure of each of the patent documents and scientific articles referred to  
5 herein is incorporated by reference for all purposes.


#### EQUIVALENTS

The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting on the invention described herein. Scope of the  
10 invention is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.


—*sig*—  
Δ<sup>2</sup>

**WHAT IS CLAIMED IS:**

1. A compound having the formula:




OR



or pharmaceutically acceptable salt, ester or prodrug thereof,

wherein:

D-Het is selected from the group consisting of:



A is selected from the group consisting of:

- a) carbonyl, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl d) -C(O)-C<sub>1-6</sub> alkyl, and  
e) -C(O)-C<sub>2-6</sub> alkenyl,

wherein

- i) 0-2 carbon atoms of the C<sub>1-6</sub> alkyl and C<sub>2-6</sub> alkenyl groups in any  
of b) – e) optionally are replaced by a moiety selected from the  
group consisting of O, S(O)<sub>p</sub>, and NR<sup>11</sup>, and  
ii) any of b) – e) optionally is substituted with one or more R<sup>12</sup>  
groups;

B is selected from the group consisting of:

- a) -C(O)NH-, b) -C(S)NH-, c) -NHC(O)-, d) -NHC(S)-, e) -S(O)<sub>2</sub>NH-,  
f) -NHS(O)<sub>2</sub>-, g) -OC(O)NH-, h) -OC(S)NH-, i) -NHC(O)NH-, j) -NHC(S)NH-,  
k) -NHC(O)O-, l) -NHC(S)O-, and m) -NR<sup>11</sup>-;


n is 0 or 1;

D is selected from the group consisting of:


- a) -CH<sub>2</sub>-, b) -C(O)-, c) -C(S)-, d) -C(=NOR<sup>11</sup>)-, e) -CH<sub>2</sub>CH<sub>2</sub>-, f) -OCH<sub>2</sub>-,  
g) -SCH<sub>2</sub>-, h) -S(O)CH<sub>2</sub>-, i) -S(O)<sub>2</sub>CH<sub>2</sub>-, j) -NR<sup>11</sup>CH<sub>2</sub>-, k) -C(O)CH<sub>2</sub>-,  
l) -C(S)CH<sub>2</sub>-, and m) -C(=NOR<sup>11</sup>)CH<sub>2</sub>-;

E is selected from the group consisting of:


a)



b)



c)



d) 5-10 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally substituted with one or more  $R^{12}$  groups;

e)  $C_{5-10}$  saturated, unsaturated, or aromatic carbocycle, optionally substituted with one or more  $R^{12}$  groups;

f)  $C_{1-8}$  alkyl,

g)  $C_{2-8}$  alkenyl,

h)  $C_{2-8}$  alkynyl,

i)  $C_{1-8}$  alkoxy,

j)  $C_{1-8}$  alkylthio,

k)  $C_{1-8}$  acyl,

l)  $S(O)_r R^{11}$ ; and

m) hydrogen,

wherein any of f) – k) optionally is substituted with

i) one or more  $R^{12}$  groups;

ii) 5-6 membered saturated, unsaturated, or aromatic heterocycle containing one or more heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally substituted with one or more  $R^{12}$  groups; or

iii)  $C_{5-10}$  saturated, unsaturated, or aromatic carbocycle, optionally substituted with one or more  $R^{12}$  groups;

M is selected from the group consisting of:

- a)  $-C(O)-$ , b)  $-C(=NOR^{11})-$ , c)  $-CH(OR^{11})-$ , d)  $-NR^{11}-CH_2-$ , e)  $-CH_2-NR^{11}-$ ,  
 f)  $-CH(NR^{11}R^{11})-$ , g)  $-C(=NNR^{11}R^{11})-$ , h)  $-NR^{11}-C(O)-$ , i)  $-C(O)NR^{11}-$ , and  
 j)  $-C(=NR^{11})-$ ;

R is selected from the group consisting of H and  $C_{1-6}$  alkyl;

$R^1$  is selected from the group consisting of:




- a) H, b) Cl, c) F, d) Br, e) I, f)  $-NR^{11}R^{11}$  g)  $-NR^{11}C(O)R^{11}$ , h)  $-OR^{11}$ ,  
 i)  $-OC(O)R^{11}$ , j)  $-OC(O)OR^{11}$ , k)  $-OC(O)NR^{11}R^{11}$ , l)  $-O-C_{1-6}$  alkyl- $R^{12}$ ,  
 m)  $-OC(O)-C_{1-6}$  alkyl- $R^{12}$ , n)  $-OC(O)O-C_{1-6}$  alkyl- $R^{12}$ ,  
 o)  $-OC(O)NR^{11}-C_{1-6}$  alkyl- $R^{12}$ , p)  $C_{1-6}$  alkyl, q)  $C_{1-6}$  alkenyl, r)  $C_{1-6}$  alkynyl,  
 wherein any of l) – r) optionally is substituted with one or more  $R^{12}$   
 groups;

$R^2$  is H;

$R^3$  is selected from the group consisting of:

- a) H, b)  $-OR^{11}$ , c)  $-O-C_{1-6}$  alkyl- $R^{12}$ , d)  $-OC(O)R^{11}$ , e)  $-OC(O)-C_{1-6}$  alkyl- $R^{12}$ ,  
 f)  $-OC(O)OR^{11}$ , g)  $-OC(O)O-C_{1-6}$  alkyl- $R^{12}$ , h)  $-OC(O)NR^{11}R^{11}$ ,  
 i)  $-OC(O)NR^{11}-C_{1-6}$  alkyl- $R^{12}$ , and  
 j)




alternatively,  $R^2$  and  $R^3$  taken together form a carbonyl group;

$R^4$  is selected from the group consisting of:

- a) H, b)  $R^{11}$ , c)  $-C(O)R^{11}$ , d)  $-C(O)OR^{11}$ , e)  $-C(O)NR^{11}R^{11}$ , f)  $-C_{1-6}$  alkyl- $G-R^{11}$ ,  
 g)  $-C_{2-6}$  alkenyl- $G-R^{11}$ , and h)  $-C_{2-6}$  alkynyl- $G-R^{11}$ ;

alternatively  $R^3$  and  $R^4$ , taken together with the atoms to which they are bonded, form:




G is selected from the group consisting of:

- a)  $-C(O)-$ , b)  $-C(O)O-$ , c)  $-C(O)NR^{11}-$ , d)  $-C(=NR^{11})-$ , e)  $-C(=NR^{11})O-$ ,  
 f)  $-C(=NR^{11})NR^{11}-$ , g)  $-OC(O)-$ , h)  $-OC(O)O-$ , i)  $-OC(O)NR^{11}-$ , j)  $-NR^{11}C(O)-$ ,  
 k)  $-NR^{11}C(O)O-$ , l)  $-NR^{11}C(O)NR^{11}-$ , m)  $-NR^{11}C(=NR^{11})NR^{11}-$ , and o)  $-S(O)_p-$ ;

$R^5$  is selected from the group consisting of:

- a)  $R^{11}$ , b)  $-OR^{11}$ , c)  $-NR^{11}R^{11}$ , d)  $-O-C_{1-6}$  alkyl- $R^{12}$ , e)  $-C(O)-R^{11}$ ,  
 f)  $-C(O)-C_{1-6}$  alkyl- $R^{12}$ , g)  $-OC(O)-R^{11}$ , h)  $-OC(O)-C_{1-6}$  alkyl- $R^{12}$ ,  
 i)  $-OC(O)O-R^{11}$ , j)  $-OC(O)O-C_{1-6}$  alkyl- $R^{12}$ , k)  $-OC(O)NR^{11}R^{11}$ ,  
 l)  $-OC(O)NR^{11}-C_{1-6}$  alkyl- $R^{12}$ , m)  $-C(O)-C_{2-6}$  alkenyl- $R^{12}$ , and  
 n)  $-C(O)-C_{2-6}$  alkynyl- $R^{12}$ ;

alternatively,  $R^4$  and  $R^5$ , taken together with the atoms to which they are bonded, form:



wherein

Q is CH or N, and

$R^{23}$  is  $-OR^{11}$ , or  $R^{11}$ ;


$R^6$  is selected from the group consisting of:

- a)  $-OR^{11}$ , b)  $-C_{1-6}$  alkoxy- $R^{12}$ , c)  $-C(O)R^{11}$ , d)  $-OC(O)R^{11}$ , e)  $-OC(O)OR^{11}$ ,  
 f)  $-OC(O)NR^{11}R^{11}$ , and g)  $-NR^{11}R^{11}$ ;

alternatively,  $R^5$  and  $R^6$  taken together with the atoms to which they are attached form a 5-membered ring by attachment to each other through a linker selected from the group consisting of:

- a)  $-OC(R^{12})_2O-$ , b)  $-OC(O)O-$ , c)  $-OC(O)NR^{11}-$ , d)  $-NR^{11}C(O)O-$ ,  
 e)  $-OC(O)NOR^{11}-$ , f)  $-NOR^{11}-C(O)O-$ , g)  $-OC(O)NNR^{11}R^{11}-$ ,  
 h)  $-NNR^{11}R^{11}-C(O)O-$ , i)  $-OC(O)C(R^{12})_2-$ , j)  $-C(R^{12})_2C(O)O-$ , k)  $-OC(S)O-$ ,  
 l)  $-OC(S)NR^{11}-$ , m)  $-NR^{11}C(S)O-$ , n)  $-OC(S)NOR^{11}-$ , o)  $-NOR^{11}-C(S)O-$ ,  
 p)  $-OC(S)NNR^{11}R^{11}-$ , q)  $-NNR^{11}R^{11}-C(S)O-$ , r)  $-OC(S)C(R^{12})_2-$ , and  
 s)  $-C(R^{12})_2C(S)O-$ ;

alternatively, M,  $R^5$ , and  $R^6$  taken together with the atoms to which they are attached form:



109

110

wherein J is selected from the group consisting of O and NR<sup>11</sup>;

111

R6' is selected from the group consisting of

112

a) -H, b) -C<sub>1-4</sub> alkyl, c) C<sub>2-4</sub> alkenyl, which can be further substituted with C<sub>1-12</sub>

113

alkyl or one or more halogens, d) C<sub>2-4</sub> alkynyl, which can be further substituted

114

with C<sub>1-12</sub> alkyl or one or more halogens, e) aryl or heteroaryl, which can be

115

further substituted with C<sub>1-12</sub> alkyl or one or more halogens, f) -C(O)H, g) -

116

COOH, h) -CN, i) -COOR<sup>11</sup>, j) -C(O)NR<sup>11</sup>R<sup>11</sup>, k) -C(O)R<sup>11</sup>, and l) -C(O)SR<sup>11</sup>,

117

wherein b) is further substituted with one or more substituents selected from the

118

group consisting of aa) -OR<sup>11</sup>, bb) halogen, cc) -SR<sup>11</sup>, dd) C<sub>1-12</sub> alkyl, which can

119

be further substituted with halogen, hydroxyl, C<sub>1-6</sub> alkoxy, or amino, ee) -OR<sup>11</sup>,

120

ff) -SR<sup>11</sup>, gg) -NR<sup>11</sup>R<sup>11</sup>, hh) -CN, ii) -NO<sub>2</sub>, jj) -NC(O)R<sup>11</sup>, kk) -COOR<sup>11</sup>, ll) -N<sub>3</sub>,

121

mm) =N-O-R<sup>11</sup>, nn) =NR<sup>11</sup>, oo) =N-NR<sup>11</sup>R<sup>11</sup>, pp) =N-NH-C(O)R<sup>11</sup>, and qq) =N-

122

NH-C(O)NR<sup>11</sup>R<sup>11</sup>;

123

alternatively R6 and R6' are taken together with the atom to which they are attached to

124

form an epoxide, a carbonyl, an olefin, or a substituted olefin, or a C<sub>3</sub>-C<sub>7</sub> carbocyclic, carbonate,

125

or carbamate, wherein the nitrogen of said carbamate can be further substituted with a C<sub>1</sub>-C<sub>6</sub>

126

alkyl;

127

R<sup>7</sup> is selected from the group consisting of:

128

a) C<sub>1-6</sub> alkyl, b) C<sub>2-6</sub> alkenyl, and c) C<sub>2-6</sub> alkynyl,

129

wherein any of a) - c) optionally is substituted with one or more R<sup>12</sup>

130

groups;

131

R<sup>8</sup> is selected from the group consisting of H and -C(O)R<sup>11</sup>;

132

R<sup>9</sup> is selected from the group consisting of H, OH, and OR<sup>11</sup>;

133

R<sup>10</sup> is selected from the group consisting of:

134

a) H, b) R<sup>11</sup>, c) -C<sub>1-6</sub> alkyl-G-R<sup>12</sup>, d) -C<sub>2-6</sub> alkenyl-G-R<sup>12</sup>, and

135

e) -C<sub>2-6</sub> alkynyl-G-R<sup>12</sup>,

136 wherein the C<sub>1-6</sub>-alkyl, C<sub>2-6</sub> alkenyl, and C<sub>2-6</sub> alkynyl group in any of  
137 c) - e) optionally is substituted with one or more R<sup>12</sup> groups;

138 R<sup>11</sup>, at each occurrence, independently is selected from the group consisting of:

139 a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>6-10</sub> saturated, unsaturated,  
140 or aromatic carbocycle, f) 3-12 membered saturated, unsaturated, or aromatic  
141 heterocycle containing one or more heteroatoms selected from the group  
142 consisting of nitrogen, oxygen, and sulfur, g) -C(O)-C<sub>1-6</sub> alkyl,  
143 h) -C(O)-C<sub>2-6</sub> alkenyl, i) -C(O)-C<sub>2-6</sub> alkynyl, j) -C(O)-C<sub>6-10</sub> saturated, unsaturated,  
144 or aromatic carbocycle, k) -C(O)-3-12 membered saturated, unsaturated, or  
145 aromatic heterocycle containing one or more heteroatoms selected from the group  
146 consisting of nitrogen, oxygen, sulfur, l) -C(O)O-C<sub>1-6</sub> alkyl,  
147 m) -C(O)O-C<sub>2-6</sub> alkenyl, n) -C(O)O-C<sub>2-6</sub> alkynyl, o) -C(O)O-C<sub>6-10</sub> saturated,  
148 unsaturated, or aromatic carbocycle, p) -C(O)O-3-12 membered saturated,  
149 unsaturated, or aromatic heterocycle containing one or more heteroatoms selected  
150 from the group consisting of nitrogen, oxygen, and sulfur, and q) -  
151 C(O)NR<sup>13</sup>R<sup>13</sup>,

152 wherein any of b) - p) optionally is substituted with one or more R<sup>12</sup>  
153 groups,

154 alternatively, NR<sup>11</sup>R<sup>11</sup> forms a 3-7 membered saturated, unsaturated or aromatic ring  
155 including the nitrogen atom to which the R<sup>11</sup> groups are bonded and optionally one or more  
156 moieties selected from the group consisting of: O, S(O)<sub>p</sub>, and NR<sup>15</sup>;

157 R<sup>12</sup> is selected from the group consisting of:

158 a) R<sup>14</sup>, b) C<sub>1-8</sub> alkyl, c) C<sub>2-8</sub> alkenyl, d) C<sub>2-8</sub> alkynyl, e) C<sub>3-12</sub> saturated,  
159 unsaturated, or aromatic carbocycle, f) 3-12 membered saturated, unsaturated, or  
160 aromatic heterocycle containing one or more heteroatoms selected from the group  
161 consisting of nitrogen, oxygen, and sulfur, and g) -NR<sup>15</sup>C(O)OR<sup>15</sup>,

162 wherein any of b) - f) optionally is substituted with one or more R<sup>14</sup>  
163 groups;

164 R<sup>13</sup>, at each occurrence, independently is selected from the group consisting of:

165 a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>3-10</sub> saturated, unsaturated,  
166 or aromatic carbocycle, and f) 3-10 membered saturated, unsaturated, or aromatic

167 heterocycle containing one or more heteroatoms selected from the group  
 168 consisting of nitrogen, oxygen, and sulfur,  
 169 wherein any of b) – f) optionally is substituted with one or more moieties selected from the  
 170 group consisting of:

171 carbonyl; formyl; F; Cl; Br; I; CN; NO<sub>2</sub>; OR<sup>15</sup>; -S(O)<sub>p</sub>R<sup>15</sup>;  
 172 -C(O)R<sup>15</sup>; -C(O)OR<sup>15</sup>; -OC(O)R<sup>15</sup>; -C(O)NR<sup>15</sup>R<sup>15</sup>;  
 173 -OC(O)NR<sup>15</sup>R<sup>15</sup>; -C(=NR<sup>15</sup>)R<sup>15</sup>; -C(R<sup>15</sup>)(R<sup>15</sup>)OR<sup>15</sup>;  
 174 -C(R<sup>15</sup>)<sub>2</sub>OC(O)R<sup>15</sup>; -C(R<sup>15</sup>)(OR<sup>15</sup>)(CH<sub>2</sub>)<sub>r</sub>NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>R<sup>15</sup>;  
 175 -NR<sup>15</sup>OR<sup>15</sup>; -NR<sup>15</sup>C(O)R<sup>15</sup>; -NR<sup>15</sup>C(O)OR<sup>15</sup>; -NR<sup>15</sup>C(O)NR<sup>15</sup>R<sup>15</sup>;  
 176 -NR<sup>15</sup>S(O)<sub>r</sub>R<sup>15</sup>; -C(OR<sup>15</sup>)(OR<sup>15</sup>)R<sup>15</sup>; -C(R<sup>15</sup>)<sub>2</sub>NR<sup>15</sup>R<sup>15</sup>; =NR<sup>15</sup>;  
 177 -C(S)NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>C(S)R<sup>15</sup>; -OC(S)NR<sup>15</sup>R<sup>15</sup>; -NR<sup>15</sup>C(S)OR<sup>15</sup>;  
 178 -NR<sup>15</sup>C(S)NR<sup>15</sup>R<sup>15</sup>; -SC(O)R<sup>15</sup>; C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl;  
 179 C<sub>2-8</sub> alkynyl; C<sub>1-8</sub> alkoxy; C<sub>1-8</sub> alkylthio; C<sub>1-8</sub> acyl; saturated,  
 180 unsaturated, or aromatic C<sub>3-10</sub> carbocycle; and saturated,  
 181 unsaturated, or aromatic 3-10 membered heterocycle containing  
 182 one or more heteroatoms selected from the group consisting of  
 183 nitrogen, oxygen, and sulfur,

184 alternatively, NR<sup>13</sup>R<sup>13</sup> forms a 3-10 membered saturated, unsaturated or aromatic ring  
 185 including the nitrogen atom to which the R<sup>13</sup> groups are attached and optionally one or more  
 186 moieties selected from the group consisting of O, S(O)<sub>p</sub>, NR<sup>15</sup>, and N;

187 alternatively, CR<sup>13</sup>R<sup>13</sup> forms a carbonyl group;

188 R<sup>14</sup>, at each occurrence, is selected from the group consisting of:

189 a) H, b) carbonyl, c) F, d) Cl, e) Br, f) I, g) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CF<sub>3</sub>, h) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>CN,  
 190 i) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NO<sub>2</sub>, j) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, k) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OR<sup>16</sup>,  
 191 l) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>S(O)<sub>p</sub>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, m) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 192 n) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>OC(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, o) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>SC(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 193 p) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)O(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, q) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>NR<sup>13</sup>C(O)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 194 r) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(O)NR<sup>13</sup>(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>, s) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NR<sup>13</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,  
 195 t) (CR<sup>13</sup>R<sup>13</sup>)<sub>r</sub>C(=NNR<sup>13</sup>R<sup>13</sup>)(CR<sup>13</sup>R<sup>13</sup>)<sub>t</sub>R<sup>16</sup>,

- 196 u)  $(\text{CR}^{13}\text{R}^{13})_r\text{C}(=\text{NNR}^{13}\text{C}(\text{O})\text{R}^{13})(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 197 v)  $(\text{CR}^{13}\text{R}^{13})_r\text{C}(=\text{NOR}^{16})(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 198 w)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{C}(\text{O})\text{O}(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 199 x)  $(\text{CR}^{13}\text{R}^{13})_r\text{OC}(\text{O})\text{NR}^{13}(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 200 y)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{C}(\text{O})\text{NR}^{13}(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 201 z)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{S}(\text{O})_p(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ , aa)  $(\text{CR}^{13}\text{R}^{13})_r\text{S}(\text{O})_p\text{NR}^{13}(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ ,  
 202 bb)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{S}(\text{O})_p\text{NR}^{13}(\text{CR}^{13}\text{R}^{13})_t\text{R}^{16}$ , cc)  $(\text{CR}^{13}\text{R}^{13})_r\text{NR}^{13}\text{R}^{13}$ ,  
 203 dd) C<sub>1-6</sub> alkyl, ee) C<sub>2-6</sub> alkenyl, ff) C<sub>2-6</sub> alkynyl, gg)  $(\text{CR}^{13}\text{R}^{13})_r\text{-C}_{3-10}$  saturated,  
 204 unsaturated, or aromatic carbocycle, and hh)  $(\text{CR}^{13}\text{R}^{13})_r\text{-3-10}$  membered  
 205 saturated, unsaturated, or aromatic heterocycle containing one or more  
 206 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,  
 207 wherein any of dd) – hh) optionally is substituted with one or more R<sup>16</sup>  
 208 groups;  
 209 alternatively, two R<sup>14</sup> groups may form -O(CH<sub>2</sub>)<sub>s</sub>O-;  
 210 R<sup>15</sup> is selected from the group consisting of:  
 211 a) H, b) C<sub>1-6</sub> alkyl, c) C<sub>2-6</sub> alkenyl, d) C<sub>2-6</sub> alkynyl, e) C<sub>3-10</sub> saturated,  
 212 unsaturated, or aromatic carbocycle, f) 3-10 membered saturated, unsaturated, or  
 213 aromatic heterocycle containing one or more heteroatoms selected from the group  
 214 consisting of nitrogen, oxygen, and sulfur, g) -C(O)-C<sub>1-6</sub> alkyl,  
 215 h) -C(O)-C<sub>1-6</sub> alkenyl, g) -C(O)-C<sub>1-6</sub> alkynyl, i) -C(O)-C<sub>3-10</sub> saturated,  
 216 unsaturated, or aromatic carbocycle, and j) -C(O)-3-10 membered saturated,  
 217 unsaturated, or aromatic heterocycle containing one or more heteroatoms selected  
 218 from the group consisting of nitrogen, oxygen, and sulfur,  
 219 wherein any of b) – j) optionally is substituted with one or more moieties  
 220 selected from the group consisting of H; F; Cl; Br; I; CN; NO<sub>2</sub>; OH; NH<sub>2</sub>;  
 221 NH(C<sub>1-6</sub> alkyl); N(C<sub>1-6</sub> alkyl)<sub>2</sub>; C<sub>1-6</sub> alkoxy; aryl; substituted aryl;  
 222 heteroaryl; substituted heteroaryl; and C<sub>1-6</sub> alkyl, optionally substituted  
 223 with one or more moieties selected from the group consisting of aryl,  
 224 substituted aryl, heteroaryl, substituted heteroaryl, F, Cl, Br, I, CN, NO<sub>2</sub>,  
 225 and OH;

226  $R^{16}$ , at each occurrence, independently is selected from the group consisting of:  
 227 a)  $R^{17}$ , b)  $C_{1-6}$  alkyl, c)  $C_{2-6}$  alkenyl, d)  $C_{2-6}$  alkynyl, e)  $-C_{3-10}$  saturated,  
 228 unsaturated, or aromatic carbocycle, and f)  $-3-10$  membered saturated,  
 229 unsaturated, or aromatic heterocycle containing one or more heteroatoms selected  
 230 from the group consisting of nitrogen, oxygen, and sulfur,  
 231 wherein any of b) – f) optionally is substituted with one or more  $R^{17}$   
 232 groups;

233  $R^{17}$ , at each occurrence, independently is selected from the group consisting of:  
 234 a) H, b) carbonyl, c) F, d) Cl, e) Br, f) I, g)  $(CR^{13}R^{13})_rCF_3$ , h)  $(CR^{13}R^{13})_rCN$ ,  
 235 i)  $(CR^{13}R^{13})_rNO_2$ , j)  $(CR^{13}R^{13})_r(CR^{13}R^{13})$ , k)  $(CR^{13}R^{13})_rOR^{11}$ ,  
 236 l)  $(CR^{13}R^{13})_rS(O)_pR^{13}$ , m)  $(CR^{13}R^{13})_rC(O)R^{13}$ , n)  $(CR^{13}R^{13})_rC(O)OR^{13}$ ,  
 237 o)  $(CR^{13}R^{13})_rOC(O)R^{13}$ , p)  $(CR^{13}R^{13})_rNR^{13}C(O)R^{13}$ ,  
 238 q)  $(CR^{13}R^{13})_rC(O)NR^{13}R^{13}$ , r)  $(CR^{13}R^{13})_rC(=NR^{13})R^{13}$ ,  
 239 s)  $(CR^{13}R^{13})_rNR^{13}C(O)NR^{13}R^{13}$ , t)  $(CR^{13}R^{13})_rNR^{13}S(O)_pR^{13}$ ,  
 240 u)  $(CR^{13}R^{13})_rS(O)_pNR^{13}R^{13}$ , v)  $(CR^{13}R^{13})_rNR^{13}S(O)_pNR^{13}R^{13}$ , w)  $C_{1-6}$  alkyl,  
 241 x)  $C_{2-6}$  alkenyl, y)  $C_{2-6}$  alkynyl, z)  $(CR^{13}R^{13})_r-C_{3-10}$  saturated, unsaturated, or  
 242 aromatic carbocycle, and aa)  $(CR^{13}R^{13})_r-3-10$  membered saturated, unsaturated,  
 243 or aromatic heterocycle containing one or more heteroatoms selected from the  
 244 group consisting of nitrogen, oxygen, and sulfur,  
 245 wherein any of w) – aa) optionally is substituted with one or more  
 246 moieties selected from the group consisting of  $R^{13}$ ; F; Cl; Br; I; CN;  $NO_2$ ;  
 247  $-OR^{13}$ ;  $-NH_2$ ;  $-NH(C_{1-6} \text{ alkyl})$ ;  $-N(C_{1-6} \text{ alkyl})_2$ ;  $C_{1-6}$  alkoxy;  $C_{1-6}$  alkylthio;  
 248 and  $C_{1-6}$  acyl;

249  $R^{18}$ , at each occurrence, independently is selected from the group consisting of:  
 250 a) H, b)  $OR^{15}$ , c)  $-O-C_{1-6} \text{ alkyl}-OC(O)R^{15}$ , d)  $-O-C_{1-6} \text{ alkyl}-OC(O)OR^{15}$ ,  
 251 e)  $-O-C_{1-6} \text{ alkyl}-OC(O)NR^{15}R^{15}$ , f)  $-O-C_{1-6} \text{ alkyl}-C(O)NR^{15}R^{15}$ ,  
 252 g)  $-O-C_{1-6} \text{ alkyl}-NR^{15}C(O)R^{15}$ , h)  $-O-C_{1-6} \text{ alkyl}-NR^{15}C(O)OR^{15}$ ,  
 253 i)  $-O-C_{1-6} \text{ alkyl}-NR^{15}C(O)NR^{15}R^{15}$ , j)  $-O-C_{1-6} \text{ alkyl}-NR^{15}C(=NH)NR^{15}R^{15}$ ,  
 254 k)  $-O-C_{1-6} \text{ alkyl}-S(O)_pR^{15}$ , l)  $-O-C_{2-6} \text{ alkenyl}-OC(O)R^{15}$ ,  
 255 m)  $-O-C_{2-6} \text{ alkenyl}-OC(O)OR^{15}$ , n)  $-O-C_{2-6} \text{ alkenyl}-OC(O)NR^{15}R^{15}$ ,

256 o) -O-C<sub>2-6</sub> alkenyl-C(O)NR<sup>15</sup>R<sup>15</sup>, p) -O-C<sub>2-6</sub> alkenyl-NR<sup>15</sup>C(O)R<sup>15</sup>,  
257 q) -O-C<sub>2-6</sub> alkenyl-NR<sup>15</sup>C(O)OR<sup>15</sup>, r) -O-C<sub>2-6</sub> alkenyl-NR<sup>15</sup>C(O)NR<sup>15</sup>R<sup>15</sup>,  
258 s) -O-C<sub>2-6</sub> alkenyl-NR<sup>15</sup>C(=NH)NR<sup>15</sup>R<sup>15</sup>, t) -O-C<sub>2-6</sub> alkenyl-S(O)<sub>p</sub>R<sup>15</sup>,  
259 u) -O-C<sub>2-6</sub> alkynyl-OC(O)R<sup>15</sup>, v) -O-C<sub>2-6</sub> alkynyl-OC(O)OR<sup>15</sup>,  
260 w) -O-C<sub>2-6</sub> alkynyl-OC(O)NR<sup>15</sup>R<sup>15</sup>, x) -O-C<sub>2-6</sub> alkynyl-C(O)NR<sup>15</sup>R<sup>15</sup>,  
261 y) -O-C<sub>2-6</sub> alkynyl-NR<sup>15</sup>C(O)R<sup>15</sup>, z) -O-C<sub>2-6</sub> alkynyl-NR<sup>15</sup>C(O)OR<sup>15</sup>,  
262 aa) -O-C<sub>2-6</sub> alkynyl-NR<sup>15</sup>C(O)NR<sup>15</sup>R<sup>15</sup>,  
263 bb) -O-C<sub>2-6</sub> alkynyl-NR<sup>15</sup>C(=NH)NR<sup>15</sup>R<sup>15</sup>, cc) -O-C<sub>2-6</sub> alkynyl-S(O)<sub>p</sub>R<sup>15</sup>; and  
264 dd) -NR<sup>15</sup>R<sup>15</sup>;

265 alternatively, two R<sup>18</sup> groups taken together form =O, =NOR<sup>15</sup>, or =NNR<sup>15</sup>R<sup>15</sup>;

266 R<sup>19</sup> is R<sup>12</sup>;

267 R<sup>20</sup> is selected from the group consisting of:

268 a) R<sup>13</sup>, b) F, c) Cl, d) Br, e) I, f) CN, g) NO<sub>2</sub>, and h) -OR<sup>11</sup>;

269 alternatively, R<sup>19</sup> and R<sup>20</sup> taken together are -O(CH<sub>2</sub>)<sub>r</sub>O-;

270 R<sup>21</sup>, at each occurrence, independently is selected from the group consisting of:

271 a) H, b) F, c) Cl, d) Br, e) I, f) CN, g) -OR<sup>11</sup>, h) NO<sub>2</sub>, i) -NR<sup>11</sup>R<sup>11</sup>, j) C<sub>1-6</sub> alkyl,

272 k) C<sub>1-6</sub> acyl, and l) C<sub>1-6</sub> alkoxy;

273 R<sup>22</sup> is selected from the group consisting of:

274 a) C<sub>1-6</sub> alkyl, b) C<sub>2-6</sub> alkenyl, c) C<sub>2-6</sub> alkynyl, d) C<sub>1-6</sub> acyl, e) C<sub>1-6</sub> alkoxy,

275 f) C<sub>1-6</sub> alkylthio, g) saturated, unsaturated, or aromatic C<sub>5-10</sub> carbocycle,

276 h) saturated, unsaturated, or aromatic 5-10 membered heterocycle containing one

277 or more heteroatoms selected from the group consisting of nitrogen, oxygen, and

278 sulfur, i) -O-C<sub>1-6</sub> alkyl-saturated, unsaturated, or aromatic 5-10 membered

279 heterocycle containing one or more heteroatoms selected from the group

280 consisting of nitrogen, oxygen, and sulfur, j) -NR<sup>11</sup>-C<sub>1-6</sub> alkyl-saturated,

281 unsaturated, or aromatic 5-10 membered heterocycle containing one or more

282 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur,

283 k) saturated, unsaturated, or aromatic 10-membered bicyclic ring system

284 optionally containing one or more heteroatoms selected from the group consisting

285 of nitrogen, oxygen, and sulfur, l) saturated, unsaturated, or aromatic 13-

286 membered tricyclic ring system optionally containing one or more heteroatoms



287 selected from the group consisting of nitrogen, oxygen, and sulfur, m) -OR<sup>11</sup>,  
288 n) -NR<sup>11</sup>R<sup>11</sup>, o) S(O)<sub>r</sub>R<sup>11</sup>, and p) R<sup>21</sup>,

289 wherein any of a) - l) optionally is substituted with one or more R<sup>12</sup>  
290 groups;

291 alternatively, R<sup>22</sup> and one R<sup>21</sup> group, taken together with the atoms to which they are  
292 bonded, form a 5-7 membered saturated or unsaturated carbocycle, optionally substituted with  
293 one or more R<sup>12</sup> groups; or a 5-7 membered saturated or unsaturated heterocycle containing one  
294 or more atoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally  
295 substituted with one or more R<sup>12</sup> groups;

296 R<sup>23</sup> at each occurrence, independently is selected from the group consisting of:

297 a) hydrogen; b) an electron-withdrawing group; c) aryl; d) substituted aryl;  
298 e) heteroaryl; f) substituted heteroaryl; and g) C<sub>1-6</sub> alkyl, optionally substituted  
299 with one or more R<sup>12</sup> groups;

300 alternatively, any R<sup>23</sup> and any R<sup>20</sup>, taken together with the atoms to which they are  
301 bonded, form a 5-7 membered saturated or unsaturated carbocycle, optionally substituted with  
302 one or more R<sup>12</sup> groups; or a 5-7 membered saturated or unsaturated heterocycle containing one  
303 or more atoms selected from the group consisting of nitrogen, oxygen, and sulfur, and optionally  
304 substituted with one or more R<sup>12</sup> groups;

305 p, at each occurrence, is selected from the group consisting of 0, 1, and 2;

306 r, at each occurrence, is selected from the group consisting of 0, 1, and 2;




307 s, at each occurrence, is selected from the group consisting of 1, 2, 3, or 4;


308 t, at each occurrence, is selected from the group consisting of 0, 1, or 2;

309 u, at each occurrence, is selected from the group consisting of 1, 2, 3, 4, or 5; and,

310 v, at each occurrence, is selected from the group consisting of 0, 1, 2, or 3.

1 2. A compound having the formula selected from the group consisting of:






5


6 or a pharmaceutically acceptable salt, ester, or prodrug thereof,

7 wherein A, B, n, D, E, R, R<sup>1</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6'</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, and R<sup>10</sup> are as defined in claim 1.


1 3. A compound having the formula selected from the group consisting of:




2



3




, and




6 or a pharmaceutically acceptable salt, ester, or prodrug thereof,

7 wherein A, B, n, D, E, R, R<sup>1</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6'</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, and R<sup>10</sup> are as defined in claim 1.


1 4. A compound having the formula selected from the group consisting of:




3




4




5





6


, and



7


8 or a pharmaceutically acceptable salt, ester, or prodrug thereof,  
 9 wherein A, B, n, E, R<sup>4</sup>, and R<sup>10</sup> are as defined in claim 1.

1 5. A compound having the formula selected from the group consisting of:




2


3




4




5



6



, and




7


8 or a pharmaceutically acceptable salt, ester, or prodrug thereof,

9 wherein A, B, n, E, R<sup>4</sup>, and R<sup>10</sup> are as defined in claim 1.

1 6. A compound having the formula selected from the group consisting of:




2




3






4




5



6

, and




7


8 or a pharmaceutically acceptable salt, ester, or prodrug thereof,

9 wherein A, B, n, E, and R<sup>10</sup> are as defined in claim 1.


1 7. A compound having the formula selected from the group consisting of:




2




3



4




5



6

, and



7  
8 or a pharmaceutically acceptable salt, ester, or prodrug thereof,  
9 wherein A, B, n, E, and R<sup>10</sup> are as defined in claim 1.


1 8. The compound according to any of claims 1-7, wherein n is 1.

1 9. The compound according to any of claims 1-8, wherein A-(B)<sub>n</sub>-D is:  
2 A-C(O)NH-D.

1 10. The compound according to any of claims 1-8, wherein A-(B)<sub>n</sub>-D is:  
2 A-SO<sub>2</sub>NH-D.




1 11. The compound according to any of claims 1-8, wherein A-(B)<sub>n</sub>-D is:  
2 A-C(S)NH-D.

1 12. A compound having the formula






3 or a pharmaceutically acceptable salt, ester, or prodrug thereof,  
4 wherein M is a macrolide selected from the group consisting of




5






6




7




8







- 93 -



- 94 -





- 95 -




23 and m is an integer from 1-4.


1 13. A compound having the formula selected from the group consisting of:








4



5



6

7 or a pharmaceutically acceptable salt, ester, or prodrug thereof.

1 14. A pharmaceutical composition comprising a compound according to any one of claims  
2 1-13 and a pharmaceutically acceptable carrier.

1 15. A method of treating a microbial infection in a mammal comprising administering to the  
2 mammal an effective amount of a compound according to any one of claims 1-13.

- 1 16. A method of treating a fungal infection in a mammal comprising administering to the  
2 mammal an effective amount of a compound according to any one of claims 1-13.
- 1 17. A method of treating a parasitic disease in a mammal comprising administering to the  
2 mammal an effective amount of a compound according to any one of claims 1-13.
- 1 18. A method of treating a proliferative disease in a mammal comprising administering to the  
2 mammal an effective amount of a compound according to any one of claims 1-13.
- 1 19. A method of treating a viral infection in a mammal comprising administering to the  
2 mammal an effective amount of a compound according to any one of claims 1-13.
- 1 20. A method of treating an inflammatory disease in a mammal comprising administering to  
2 the mammal an effective amount of a compound according to any one of claims 1-13.
- 1 21. A method of treating a gastrointestinal motility disorder in a mammal comprising  
2 administering to the mammal an effective amount of a compound according to any one of claims  
3 1-13.
- 1 22. The method according to any one of claims 15-21 wherein the compound is administered  
2 orally, parentally, or topically.
- 1 23. A method of synthesizing a compound according to any of claims 1-13.
- 1 24. A medical device containing a compound according to any one of claims 1-13.
- 1 25. The medical device according to claim 24, wherein the device is a stent.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/036146

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H17/00 C07H17/08 A61K31/7048 A61K31/7052 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| A          | WO 03/070174 A (SYMPORE GMBH; BURNET, MICHAEL; GUSE, JAN-HINRICH; GUTKE, HANS-JURGEN;) 28 August 2003 (2003-08-28)                |                       |
| A          | WO 99/63937 A (ADVANCED MEDICINE, INC; GRIFFIN, JOHN, H; PACE, JOHN, L) 16 December 1999 (1999-12-16)<br>cited in the application |                       |
| A          | US 6 034 069 A (OR ET AL) 7 March 2000 (2000-03-07)<br>cited in the application   |                       |

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

23 March 2005

Date of mailing of the international search report

01/04/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bardili, W

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/036146

| Patent document<br>cited in search report |            | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|------------|---------------------|----------------------------|---------------------|
| WO 03070174                               | A          | 28-08-2003          | AU 2003219770 A1           | 09-09-2003          |
|   |            |                     | CA 2476423 A1              | 28-08-2003          |
|   |            |                     | EP 1483277 A2              | 08-12-2004          |
|   |            |                     | WO 03070174 A2             | 28-08-2003          |
|   |            |                     | US 2004087517 A1           | 06-05-2004          |
|   |            |                     | US 2004186063 A1           | 23-09-2004          |
| <hr/>                                     |            |                     |                            |                     |
| WO 9963937                                | A          | 16-12-1999          | AU 4426399 A               | 30-12-1999          |
|   |            |                     | AU 4426699 A               | 30-12-1999          |
|   |            |                     | AU 4428899 A               | 30-12-1999          |
|   |            |                     | AU 4430599 A               | 30-12-1999          |
|   |            |                     | AU 4543699 A               | 30-12-1999          |
|   |            |                     | AU 4543799 A               | 30-12-1999          |
|   |            |                     | AU 4543899 A               | 30-12-1999          |
|   |            |                     | AU 4551699 A               | 30-12-1999          |
|   |            |                     | AU 4551799 A               | 30-12-1999          |
|   |            |                     | AU 764401 B2               | 21-08-2003          |
|   |            |                     | AU 4551899 A               | 30-12-1999          |
|   |            |                     | AU 4674599 A               | 30-12-1999          |
|   |            |                     | AU 4677199 A               | 30-12-1999          |
|   |            |                     | AU 4677399 A               | 30-12-1999          |
|   |            |                     | AU 4818099 A               | 30-12-1999          |
|   |            |                     | CA 2316999 A1              | 16-12-1999          |
|   |            |                     | CA 2318547 A1              | 16-12-1999          |
|   |            |                     | CA 2319080 A1              | 16-12-1999          |
|   |            |                     | CA 2319120 A1              | 16-12-1999          |
|   |            |                     | CA 2319474 A1              | 16-12-1999          |
|   |            |                     | CA 2319477 A1              | 16-12-1999          |
|   |            |                     | CA 2319495 A1              | 16-12-1999          |
|   |            |                     | CA 2319761 A1              | 16-12-1999          |
|   |            |                     | CA 2320241 A1              | 16-12-1999          |
|   |            |                     | CA 2320290 A1              | 16-12-1999          |
|   |            |                     | CA 2321162 A1              | 16-12-1999          |
|   |            |                     | CA 2321166 A1              | 16-12-1999          |
|   |            |                     | CA 2321190 A1              | 16-12-1999          |
|   |            |                     | CA 2321274 A1              | 16-12-1999          |
|   |            |                     | EP 1083919 A1              | 21-03-2001          |
|   |            |                     | EP 1083920 A1              | 21-03-2001          |
|   |            |                     | EP 1085845 A2              | 28-03-2001          |
|   |            |                     | EP 1085861 A1              | 28-03-2001          |
|   |            |                     | EP 1083894 A1              | 21-03-2001          |
|   |            |                     | EP 1086065 A1              | 28-03-2001          |
|   |            |                     | EP 1085846 A2              | 28-03-2001          |
|   |            |                     | EP 1124528 A1              | 22-08-2001          |
|   |            |                     | EP 1003541 A1              | 31-05-2000          |
|   |            |                     | EP 1005356 A1              | 07-06-2000          |
|   |            |                     | EP 0989857 A1              | 05-04-2000          |
|   |            |                     | EP 1085891 A1              | 28-03-2001          |
|   |            |                     | EP 1143991 A2              | 17-10-2001          |
|   |            |                     | EP 1085893 A1              | 28-03-2001          |
|   |            |                     | JP 2002517422 T            | 18-06-2002          |
|   |            |                     | JP 2002517423 T            | 18-06-2002          |
|   |            |                     | JP 2002517458 T            | 18-06-2002          |
|   |            |                     | JP 2002517465 T            | 18-06-2002          |
| JP 2004512255 T                           | 22-04-2004 |                     |                            |                     |
| NZ 505979 A                               | 26-09-2003 |                     |                            |                     |
| SG 80631 A1                               | 22-05-2001 |                     |                            |                     |

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/036146

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| US 6034069                                | A                   | 07-03-2000                 | NONE                |

**THIS PAGE BLANK (USPTO)**